arXiv:1608.02049v1 [math.AG] 6 Aug 2016

Weighted complete intersection del Pezzo surfaces

Evgeny Mayanskiy

August 9, 2016

Abstract

We classify codimension 2 well-formed and quasi-smooth weighted complete intersection del Pezzo surfaces.

1 Introduction

According to [3], Theorem 1.3, a weighted complete intersection del Pezzo surface, which is well-formed, quasi-smooth and is not an intersection with a linear cone, has codimension 0, 1 or 2. Weighted 2-dimensional del Pezzo hypersurfaces were studied in [6], [1], [2] and others. Weighted codimension 2 complete intersection del Pezzo surfaces of amplitude 1 were classified in [7].

The goal of this note is to classify weighted complete intersection del Pezzo surfaces of codimension 2.

Our main result is the following. The reader may refer to the next section for the notation and terminology.

Main Theorem. *Let $a_0 \le a_1 \le a_2 \le a_3 \le a_4$, $d_1 \le d_2$, $X \subset \mathbb{P}(a_0, a_1, a_2, a_3, a_4)$ be a well-formed quasi-smooth weighted complete intersection del Pezzo surface, given by the intersection of two quasi-homogeneous polynomials of degrees d_1 and d_2, which is not an intersection with a linear cone. Then one of the following holds:*

(1) either $a_4 \le 500$, $d_2 \le 1000$,

(2) or $(a_0, a_1, a_2, a_3, a_4;\ d_1, d_2)$ appears in Table 1 (see Appendix).

In other words, we explicitly classify infinite series del Pezzo surface complete intersections and leave the computation of the "sporadic cases" in (1) to a computer. The latter is straightforward using Theorem WF and Theorem QS of Iano-Fletcher quoted in the next section. See Table 2 in Appendix.

2 Notation and assumptions

We work over a fixed algebraically closed field of characteristic 0. Our terminology and notation for weighted complete intersections and projective spaces follow [5].

$\mathbb{P} = \mathbb{P}(a_0, \dots, a_n)$, $a_0 \le \dots \le a_n$, denotes the weighted projective space with weights $a_0, \dots, a_n$ on the coordinates $x_0, \dots, x_n$.

A weighted complete intersection $X \subset \mathbb{P}$ of codimension c is assumed to be given by *general* quasi-homogeneous polynomials $F_1, \dots, F_c$ of degrees $d_1, \dots, d_c$, $d_1 \le \dots \le d_c$. The generality hypothesis simplifies the exposition and is not essential otherwise. It implies that a monomial $x_0^{m_0} \dots x_n^{m_n}$ appears in F_i with

a nonzero coefficient if and only if $d_i = \sum_j m_j \cdot a_j$.

We restrict our attention only to those weighted complete intersections, which are *not intersections with linear cones*, i.e. no degree d_i is equal to one of the weights $a_0, \ldots, a_n$.

Our classification is based on the following criteria proved in [5]. The reader may refer to [5] for the notions of well-formedness and quasi-smoothness.

Denote $b_{i_1 \cdots i_k} = \gcd(a_0 \ldots \hat{a}_{i_1} \ldots \hat{a}_{i_k} \ldots a_n)$, $i_1 < \ldots < i_k$. Let $X \subset \mathbb{P}(a_0, a_1, a_2, a_3, a_4)$ be a codimension $c = 2$ weighted complete intersection (general and not an intersection with a linear cone).

Theorem WF **(cf. [5],** 6.11**).** *X is well-formed if and only if*

(1) $\forall\ i < j < k,\ b_{ijk} \mid d_1$ *or* $b_{ijk} \mid d_2$,

(2) $\forall\ i < j,\ b_{ij} \mid d_1$ *and* $b_{ij} \mid d_2$,

(3) $\forall i,\ b_i = 1$.

Theorem QS **(cf. [5],** 8.7**).** *X is quasi-smooth if and only if*

(1) $\forall i \in \{0, 1, 2, 3, 4\}$,

- *either* $d_1 \vdots a_i$,
- *or* $d_2 \vdots a_i$,
- *or* $d_1 - a_e,\ d_2 - a_f \in (a_i)$ *for some* $e \neq f$;

(2) $\forall\ 0 \leq i < j \leq 4$,

- *either* $d_1, d_2 \in (a_i, a_j)$,
- *or* $d_1,\ d_2 - a_e \in (a_i, a_j)$ *for some* e,
- *or* $d_1 - a_e,\ d_2 \in (a_i, a_j)$ *for some* e,
- *or* $d_1 - a_e,\ d_2 - a_f \in (a_i, a_j),\ \ \#\{e\} = \#\{f\} = 2,\ \{e, f\} = \{k, l, m\}$;

(3) $\forall\ 0 \leq k < l < m \leq 4$,

- *either* $d_1, d_2 \in (a_k, a_l, a_m)$,
- *or* $d_1,\ d_2 - a_i,\ d_2 - a_j \in (a_k, a_l, a_m)$,
- *or* $d_1 - a_i,\ d_1 - a_j,\ d_2 \in (a_k, a_l, a_m)$.

Here we assume that $\{i, j, k, l, m\} = \{0, 1, 2, 3, 4\}$ and denote $(a_{i_1}, \ldots, a_{i_p}) = \{\sum_{s=1}^{p}\ u_s \cdot a_{i_s}\ \mid\ u_s \in \mathbb{Z},\ u_s \geq 0\}$.

The *amplitude* of X is defined as

$$I = a_0 + a_1 + a_2 + a_3 + a_4 - d_1 - d_2.$$

In accordance with the adjunction formula ([4], Theorem 3.3.4, [5], 6.14), we give the following definition.

Definition. A well-formed and quasi-smooth weighted complete intersection $X \subset \mathbb{P}(a_0, a_1, a_2, a_3, a_4)$ of dimension 2 is a *del Pezzo surface*, if

$$I \geq 1. \tag{1}$$

Throughout the note, λ, μ, ν, α, β, γ denote integers.

3 Preliminary analysis

The proof of the Main Theorem is an elementary, but somewhat lengthy analysis of all sequences (a_0, a_1, a_2, a_3, a_4; d_1, d_2), which satisfy the conditions of Theorem WF and Theorem QS of Iano-Fletcher quoted above, as well as inequality (1). In this section we list all possibilities for the expressions of d_1 and d_2 in terms of the a_i's. Each of the resulting cases is studied separately in this and the next sections. All together they lead to the proof of the Main Theorem.

Lemma 1. *Let $a_0 \le a_1 \le a_2 \le a_3 \le a_4$, $d_1 \le d_2$, $X \subset \mathbb{P}(a_0, a_1, a_2, a_3, a_4)$ be a well-formed quasi-smooth weighted complete intersection del Pezzo surface, given by the intersection of two quasi-homogeneous polynomials of degrees d_1 and d_2, which is not an intersection with a linear cone.*

Then (d_1, d_2) is one of the following:

(01) $d_1 = a_0 + a_4$, $d_2 = a_1 + a_4$; *(02)* $d_1 = a_0 + a_4$, $d_2 = a_2 + a_4$;

(12) $d_1 = a_1 + a_4$, $d_2 = a_2 + a_4$; *(03)* $d_1 = a_0 + a_4$, $d_2 = a_3 + a_4$;

(13) $d_1 = a_1 + a_4$, $d_2 = a_3 + a_4$; *(23)* $d_1 = a_2 + a_4$, $d_2 = a_3 + a_4$;

(4.1) $d_1 = a_0 + a_3$, $d_2 = 2a_4$; *(4.2)* $d_1 = a_1 + a_3$, $d_2 = 2a_4$; *(4.3)* $d_1 = a_2 + a_3$, $d_2 = 2a_4$;

(4.4) $d_1 = a_0 + a_4$, $d_2 = 2a_4$; *(4.5)* $d_1 = a_1 + a_4$, $d_2 = 2a_4$; *(4.6)* $d_1 = a_2 + a_4$, $d_2 = 2a_4$;

(4.7) $d_1 = 2a_3$, $d_2 = 2a_4$; *(4.8)* $d_1 = a_3 + a_4$, $d_2 = 2a_4$; *(4.9)* $d_1 = 2a_4$, $d_2 = 2a_4$.

Proof: First, note that Theorem QS (2), $(i, j) = (3, 4)$, implies that $d_1 \ge a_0 + a_3$.

Hence (1) gives

$$d_2 < a_1 + a_2 + a_4. \tag{2}$$

By Theorem QS (1), $i = 4$, we have:

- either $d_2 \vdots a_4$, i.e. $d_2 = 2a_4$ by (2),
- or $d_1 \vdots a_4$, i.e. $d_1 = d_2 = 2a_4$ by (1) and Theorem QS (2), $(i, j) = (3, 4)$,
- or $d_1 = a_e + \lambda a_4$, $d_2 = a_f + \mu a_4$ for some $e \ne f$, $\mu, \lambda \ge 1$.

In the last case, (1) implies that $\lambda + \mu \le 2$, i.e. $\lambda = \mu = 1$. This gives items (01), (02), (12), (03), (13), (23).

If $d_2 = 2a_4$, then Theorem QS (2), $(i, j) = (3, 4)$, requires $d_1 - a_e \in (a_3, a_4)$, which gives items (4.1)$-$(4.9). *QED*

Corollary 1. *Under the assumptions of Lemma* 1, $d_2 \le 2a_4$.

The next lemma will be used frequently.

Lemma 2. *Under the assumptions of Lemma* 1, *if* $a_0 = \lambda_0 a_i$, $a_1 = \lambda_1 a_i$, $a_2 = \lambda_2 a_i$, $a_3 = \lambda_3 a_i$, $a_4 = \lambda_4 a_i$ *for some i, then*

$$a_4 \le \lambda_4 \cdot \min_j \mathrm{lcm}(\textit{denominators of } \lambda_0, \dots, \hat{\lambda}_j, \dots, \lambda_4).$$

Proof: By Theorem WF, $a_i = \mathrm{lcm}(\text{denominators of } \lambda_0, \dots, \hat{\lambda}_j, \dots, \lambda_4)$ for any j, if λ_k's are reduced fractions. *QED*

Lemma 3. *Under the assumptions of Lemma* 1, $a_4 \le 3$, *whenever three of the weights* a_i *coincide.*

Proof: There are three cases.

Case 1. Assume $a_0 = a_1 = a_2$. Then by Theorem WF, $d_1 \vdots a_0$ and $d_2 \vdots a_0$.

If $d_2 \ge a_3 + a_4$, then $d_1 = 2a_0 \ge a_0 + a_3$, i.e. $a_0 = a_1 = a_2 = a_3$, which should be 1 by Theorem WF. Then (1) implies that $a_4 = 1$ as well.

Otherwise, by Lemma 1, $d_1 = d_2 = a_0 + a_4$, and so $a_4 \vdots a_0$. Hence by Theorem WF, $a_0 = a_1 = a_2 = 1$, $d_1 = d_2 = 1 + a_4$. By (1), $a_3 = a_4$, which should be 1 by Theorem WF.

Case 2. Assume $a_1 = a_2 = a_3$. Then by Theorem WF, $d_1 \vdots a_1$ and $d_2 \vdots a_1$. Hence by Lemma 1, $2a_4 \vdots a_1$, i.e. $a_1 \in \{1, 2\}$ by Theorem WF. Then (1) and Lemma 1 imply that $a_4 < 2a_1 \le 4$.

Case 3. Assume $a_2 = a_3 = a_4$. Then $d_1 = d_2 = 2a_4$ and (1) gives:

$$a_4 < a_0 + a_1.$$

By Theorem QS (1), $i = 1$, $2a_4 \vdots a_1$. Then $a_1 = 2$ by Theorem WF, and so $a_4 < 2a_1 = 4$. *QED*

Lemma 4. *The* Main Theorem *holds, if* $a_i = a_j$ *for some* $i \ne j$.

Proof: There are four cases. In each case we consider separately the subcases arising from Lemma 1.

Case ($a_0 = a_1$): *Subcase* (01)*:* $d_1 = d_2 = a_0 + a_4$, $a_4 < a_2 + a_3$ by (1).

By Theorem QS (1), $i = 3$,

- either $a_0 + a_4 = \mu a_3$, where $\mu = 2$ by (1), i.e. $a_4 = 2a_3 - a_0$,
- or $a_4 \vdots a_3$, and so $a_4 = a_3$ by (1).

If $a_4 = a_3$, then $d_1 = d_2 = a_0 + a_3 \vdots a_3$ by Theorem WF. This situation was considered in Lemma 3.

So, we may assume that $a_4 = 2a_3 - a_0$, $d_1 = d_2 = 2a_3$ and $a_3 < a_0 + a_2$. Then Theorem QS (1), $i = 2$, gives:

- either $2a_3 \vdots a_2$, i.e. $a_3 = a_2$ or $a_3 = 3a_2/2$ by (1),
- or $a_3 = a_2 + a_0/2$.

If $a_3 = a_2$, then $d_1 = d_2 = 2a_2 \vdots a_0$ by Theorem WF, i.e.

- either $a_0 = a_1 = 1$, $a_2 = a_3 = t$, $a_4 = 2t - 1$, $d_1 = d_2 = 2t$,
- or $a_0 = a_1 = 2$, $a_2 = a_3 = t$, $a_4 = 2t - 2$, $d_1 = d_2 = 2t$.

Note that in the second case t must be odd by Theorem WF. Both these cases appear in Table 1 (No. 15 and No. 21).

If $a_3 = 3a_2/2$, then $a_2 < 2a_0$ by (1) and $d_1 = d_2 = 3a_2 \vdots a_0$ by Theorem WF, i.e. $a_2 = \lambda a_0/3$, $a_1 = a_0$, $a_4 = (\lambda - 1)a_0$, $\lambda < 6$ by (1), and so $a_4 \le 3(\lambda - 1) < 15$ by Lemma 2.

If $a_3 = a_2 + a_0/2$, then $d_1 = d_2 = 2a_2 + a_0$. By Theorem WF, $2a_2 \vdots a_0$, and so

$$a_0 = a_1 = 2,\ a_2 = t,\ a_3 = t + 1,\ a_4 = 2t,\ d_1 = d_2 = 2t + 2.$$

Since either a_2 or a_3 is even, this situation violates conditions of Theorem WF.

Subcase (02) = (12)*:* $d_1 = a_0 + a_4$, $d_2 = a_2 + a_4$, $a_4 < a_0 + a_3$ by (1).

By Theorem QS (1), $i = 3$,

- either $a_0 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_0$ by (1),
- or $a_2 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_2$ by (1),
- or $a_0 + a_4 = a_e + \lambda a_3$ and $a_2 + a_4 = a_f + \mu a_3$ for some $e \neq f$, and so $a_4 = a_3$ or ($a_3 = a_2$, $a_4 = 2a_2 - a_0$) or ($a_3 = 2(a_2 - a_0)$, $a_4 = 3(a_2 - a_0)$).

If $a_4 = a_3$, then $a_i = 1$ for all i by Theorem WF.

If $a_2 = a_3$, $a_4 = 2a_2 - a_0$, then $d_1 = 2a_2$, $d_2 = 3a_2 - a_0$, $a_2 < 2a_0$. By Theorem WF, $6a_2 \vdots a_0$, i.e. $a_2 = a_3 = \lambda a_0/6$, $\lambda < 12$, and so $a_4 = (\lambda/3 - 1)a_0 < 24$ by Lemma 2.

If $a_3 = 2(a_2 - a_0)$, $a_4 = 3(a_2 - a_0)$, then $d_1 = a_0 + 3(a_2 - a_0)$, $d_2 = a_0 + 4(a_2 - a_0)$. Hence $a_0 \vdots a_2 - a_0$ by Theorem WF, i.e.

$$a_0 = a_1 = \frac{\lambda}{1+\lambda} a_2, \text{ and so } \quad a_4 = \frac{3}{1+\lambda} a_2 \leq 3 \quad \text{by Lemma 2.}$$

If $a_4 = 2a_3 - a_0$, then $d_1 = 2a_3$, $d_2 = 2a_3 + a_2 - a_0$, $a_3 < 2a_0$.

By Theorem QS (1), $i = 2$,

- either $2a_3 \vdots a_2$, i.e. $a_3 = a_2$ or $a_3 = 3a_2/2$ by (1),
- or $a_3 = a_2 + a_0/2$.

The case $a_2 = a_3$ was considered above. If $a_3 = 3a_2/2$, then $d_1 = 3a_2$, $d_2 = 4a_2 - a_0$, $a_2 < 4a_0/3$. By Theorem WF, $12a_2 \vdots a_0$, i.e. $a_2 = \lambda a_0/12$, $\lambda < 16$, and so $a_4 = (\lambda/4 - 1)a_0 < 48$ by Lemma 2.

If $a_3 = a_2 + a_0/2$, then $d_1 = 2a_2 + a_0$, $d_2 = 3a_2$, $a_2 < 3a_0/2$. By Theorem WF, $6a_2 \vdots a_0$, i.e. $a_2 = \lambda a_0/6$, $\lambda < 9$, and so $a_4 = (\lambda/3)a_0 < 18$ by Lemma 2.

If $a_4 = 2a_3 - a_2$, then $d_1 = 2a_3 - a_2 + a_0$, $d_2 = 2a_3$, $a_3 < a_0 + a_2$.

By Theorem QS (1), $i = 2$,

- either $2a_3 \vdots a_2$, i.e. $a_3 = a_2$ or $a_3 = 3a_2/2$ by (1),
- or $2a_3 + a_0 \vdots a_2$, i.e. $a_3 = 2a_2 - a_0/2$ or $a_3 = 3a_2/2 - a_0/2$ by (1),
- or $a_3 = 2a_2 - a_0$.

The case $a_2 = a_3 = a_4$ was considered in Lemma 3.

If $a_3 = 3a_2/2$, then $d_1 = 2a_2 + a_0$, $d_2 = 3a_2$, $a_2 < 2a_0$. By Theorem WF, $6a_2 \vdots a_0$, i.e. $a_2 = \lambda a_0/6$, $\lambda < 12$, and so $a_4 = (\lambda/3)a_0 < 24$ by Lemma 2.

If $a_3 = 2a_2 - a_0/2$, then $d_1 = 3a_2$, $d_2 = 4a_2 - a_0$, $a_2 < 3a_0/2$. By Theorem WF, $12a_2 \vdots a_0$, i.e. $a_2 = \lambda a_0/12$, $\lambda < 18$, and so $a_4 = (\lambda/4 - 1)a_0 < 54$ by Lemma 2.

If $a_3 = 3a_2/2 - a_0/2$, then $d_1 = 2a_2$, $d_2 = 3a_2 - a_0$, $a_2 < 3a_0$. By Theorem WF, $6a_2 \vdots a_0$, i.e. $a_2 = \lambda a_0/6$, $\lambda < 18$, and so $a_4 = (\lambda/3 - 1)a_0 < 36$ by Lemma 2.

If $a_3 = 2a_2 - a_0$, then $d_1 = 3a_2 - a_0$, $d_2 = 4a_2 - 2a_0$, $a_2 < 2a_0$. By Theorem WF, $12a_2 \vdots a_0$, i.e. $a_2 = \lambda a_0/12$, $\lambda < 24$, and so $a_4 = (\lambda/4 - 2)a_0 < 72$ by Lemma 2.

Subcases (03) = (13), (23), (4.1) – (4.9)*:* The same analysis as in the previous subcases gives:

$$a_4 < 60.$$

Case ($a_1 = a_2$): *Subcase* (01) = (02)*:* $d_1 = a_0 + a_4$, $d_2 = a_1 + a_4$, $a_4 < a_1 + a_3$ by (1).

By Theorem QS (1), $i = 3$,

- either $a_0 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_0$ by (1),
- or $a_1 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_1$ by (1),
- or $a_0 + a_4 = a_e + \lambda a_3$ and $a_1 + a_4 = a_f + \mu a_3$ for some $e \neq f$, and so $a_4 = a_3$ or ($a_3 = 2(a_1 - a_0)$, $a_4 = 3(a_1 - a_0)$).

If $a_3 = 2(a_1 - a_0)$, $a_4 = 3(a_1 - a_0)$, then $d_1 = a_0 + 3(a_1 - a_0)$, $d_2 = a_0 + 4(a_1 - a_0)$. Hence $a_0 \vdots a_1 - a_0$ by Theorem WF, and so

$$a_1 = a_2 = \frac{1+\lambda}{\lambda} a_0,\ a_4 = \frac{3}{\lambda} a_0 \leq 3 \quad \text{by Lemma 2.}$$

If $a_3 = a_4$, then $a_1 = a_4$ by Theorem WF. This situation was considered in Lemma 3.

If $a_4 = 2a_3 - a_1$, then $d_1 = 2a_3 + a_0 - a_1$, $d_2 = 2a_3$, $a_3 < 2a_1$. By Theorem WF,

- either $2a_3 \vdots a_1$, i.e. $a_3 = 3a_1/2$ by (1),
- or $2a_3 + a_0 \vdots a_1$, i.e. $a_3 = 2a_1 - a_0/2$ or $a_3 = 3a_1/2 - a_0/2$ by (1).

If $a_3 = 3a_1/2$, then $a_4 = 2a_1 \leq 4$ by Lemma 2.

If $a_3 = 2a_1 - a_0/2$, then $d_1 = 3a_1$, $d_2 = 4a_1 - a_0$. By Theorem QS (1), $i = 0$,

- either $3a_1 \vdots a_0$,
- or $4a_1 \vdots a_0$, i.e. $a_1 = a_2 = \lambda a_0/4$, $a_3 = (\lambda - 1)a_0/2$, $a_4 = (3\lambda/4 - 1)a_0$, $d_1 = 3\lambda a_0/4$, $d_2 = (\lambda - 1)a_0$.

If $3a_1 \vdots a_0$, then a_0, $a_1 = a_2$ and a_4 are even, which violates conditions of Theorem WF.

Hence

- either $a_0 = 2$, $a_1 = a_2 = t$, $a_3 = 2t - 1$, $a_4 = 3t - 2$, $d_1 = 3t$, $d_2 = 4t - 2$, where t is odd,
- or $a_0 = 4$, $a_1 = a_2 = 2t + 1$, $a_3 = 4t$, $a_4 = 6t - 1$, $d_1 = 6t + 3$, $d_2 = 8t$.

These solutions appear in Table 1 (No. 25 and No. 35).

If $a_3 = 3a_1/2 - a_0/2$, then $d_1 = 2a_1$, $d_2 = 3a_1 - a_0$. By Theorem QS (1), $i = 0$,

- either $2a_1 \vdots a_0$, i.e. $a_1 = a_2 = \lambda a_0/2$, $a_3 = (3\lambda - 2)a_0/4$, $a_4 = (\lambda - 1)a_0$, $d_1 = \lambda a_0$, $d_2 = (3\lambda/2 - 1)a_0$,
- or $3a_1 \vdots a_0$, i.e. $a_1 = a_2 = \lambda a_0/3$, $a_3 = (\lambda - 1)a_0/2$, $a_4 = (2\lambda/3 - 1)a_0$, $d_1 = 2\lambda a_0/3$, $d_2 = (\lambda - 1)a_0$.

Hence

- either $a_0 = 1$, $a_1 = a_2 = 2t + 1$, $a_3 = 3t + 1$, $a_4 = 4t + 1$, $d_1 = 4t + 2$, $d_2 = 6t + 2$,
- or $a_0 = 3$, $a_1 = a_2 = 2t + 1$, $a_3 = 3t$, $a_4 = 4t - 1$, $d_1 = 4t + 2$, $d_2 = 6t$.

Note that in the second case $t \not\equiv 1 \pmod 3$ by Theorem WF. These solutions appear in Table 1 (No. 19 and No. 30).

If $a_4 = 2a_3 - a_0$, then $d_1 = 2a_3$, $d_2 = 2a_3 + a_1 - a_0$, $a_3 < a_0 + a_1$. By Theorem WF,

- either $2a_3 \vdots a_1$, i.e. $a_3 = 3a_1/2$ by (1),
- or $2a_3 - a_0 \vdots a_1$, i.e. $a_3 = a_1 + a_0/2$ by (1).

If $a_3 = a_1 + a_0/2$, then $a_4 = 2a_1$, $d_1 = 2a_1 + a_0$, $d_2 = 3a_1$. By Theorem WF, $a_0 = a_1 = a_2$. This situation was considered in Lemma 3.

If $a_3 = 3a_1/2$, then $d_1 = 3a_1$, $d_2 = 4a_1 - a_0$, $a_1 < 2a_0$. By Theorem QS (1), $i = 0$, either $12a_1 \vdots a_0$ or $5a_1 \vdots a_0$. If $12a_1 \vdots a_0$, then $a_1 = a_2 = \lambda a_0/12$, $\lambda < 24$, and so $a_4 = (\lambda/4 - 1)a_0 < 60$ by Lemma 2. If $5a_1 \vdots a_0$, then $a_1 = a_2 = \lambda a_0/5$, $\lambda < 10$, and so $a_4 = (3\lambda/5 - 1)a_0 < 30$ by Lemma 2.

Subcase (12)*:* $d_1 = d_2 = a_1 + a_4$, $a_4 < a_0 + a_3$ by (1).

By Theorem QS (1), $i = 3$,

- either $a_1 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_1$ by (1),
- or $a_4 \vdots a_3$, i.e. $a_4 = a_3$ by (1).

If $a_4 = 2a_3 - a_1$, then $d_1 = d_2 = 2a_3$, $a_3 < a_0 + a_1$. By Theorem WF, $2a_3 \vdots a_1$, i.e. $a_3 = 3a_1/2$ by (1). Then $a_4 = 2a_1 \le 4$ by Lemma 2.

If $a_3 = a_4$, then $d_1 = d_2 = a_1 + a_3$. By Theorem WF, $a_3 \vdots a_1$, i.e. $a_3 = a_4 = \lambda a_1$. Hence $a_0 = a_1 = a_2 = 1$ by Lemma 2. This situation was considered in Lemma 3.

Subcase (03)*:* $d_1 = a_0 + a_4$, $d_2 = a_3 + a_4$, $a_4 < 2a_1$ by (1).

By Theorem QS (1), $i = 3$,

- either $a_4 \vdots a_3$, i.e. $a_4 = a_3$ by (1),
- or $a_0 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_0$ by (1),
- or $a_4 = a_e + \lambda a_3$ and $a_0 + a_4 = a_f + \mu a_3$ for some $e \ne f$, and so $a_4 = a_3 + a_1 - a_0$.

If $a_4 = a_3 + a_1 - a_0$, then $d_1 = a_1 + a_3$, $d_2 = 2a_3 + a_1 - a_0$, $a_3 < a_0 + a_1$. By Theorem WF,

- either $a_3 \vdots a_1$, i.e. $a_1 = a_2 = a_3$ by (1),
- or $2a_3 - a_0 \vdots a_1$, i.e. $a_3 = a_1 + a_0/2$ by (1).

If $a_3 = a_1 + a_0/2$, then $d_1 = 2a_1 + a_0/2$, $d_2 = 3a_1$. By Theorem QS (1), $i = 0$,

- either $4a_1 \vdots a_0$, i.e. $a_1 = a_2 = \lambda a_0/4$, $a_3 = (\lambda+2)a_0/4$, $a_4 = (\lambda-1)a_0/2$, $d_1 = (\lambda+1)a_0/2$, $d_2 = 3\lambda a_0/4$,
- or $3a_1 \vdots a_0$.

If $3a_1 \vdots a_0$, then a_0, $a_1 = a_2$ are even. Hence d_1 should be even by Theorem WF. This implies that $4 \mid a_0$, and so a_4 is even too. This violates conditions of Theorem WF.

If $4a_1 \vdots a_0$ and $a_0 = 2$, then a condition of Theorem WF is violated. Hence

$$a_0 = 4,\ a_1 = a_2 = 2t + 1,\ a_3 = 2t + 3,\ a_4 = 4t,\ d_1 = 4t + 4,\ d_2 = 6t + 3.$$

This solution appears in Table 1 (No. 34).

If $a_4 = 2a_3 - a_0$, then $d_1 = 2a_3$, $d_2 = 3a_3 - a_0$, $a_3 < a_1 + a_0/2$. By Theorem WF,

- either $2a_3 \vdots a_1$, i.e. $a_1 = a_2 = a_3$,
- or $3a_3 - a_0 \vdots a_1$, i.e. $a_3 = a_1 + a_0/3$.

If $a_3 = a_1 + a_0/3$, then $d_1 = 2a_1 + 2a_0/3$, $d_2 = 3a_1$. By Theorem QS (1), $i = 0$, $6a_1 \vdots a_0$, i.e.

$$a_1 = a_2 = \frac{\lambda a_0}{6},\ a_3 = \frac{\lambda+2}{6}a_0,\ a_4 = \frac{\lambda-1}{3}a_0,\ d_1 = \frac{\lambda+2}{3}a_0,\ d_2 = \frac{\lambda a_0}{2}.$$

This violates the condition of Theorem QS (2), $(i,j) = (1,2)$.

If $a_4 = a_3$, then $d_1 = a_0 + a_3$, $d_2 = 2a_3$, $a_3 < 2a_1$. By Theorem WF,

- either $2a_3 \vdots a_1$, i.e. $a_3 = 3a_1/2$,
- or $a_3 + a_0 \vdots a_1$, i.e. $a_3 = 2a_1 - a_0$.

If $a_3 = 3a_1/2$, then $a_1 = 2$ by Theorem WF, and so $a_4 = 3$. If $a_3 = 2a_1 - a_0$, then $d_1 = 2a_1$, $d_2 = 4a_1 - 2a_0$. By Theorem QS (1), $i = 0$,

- either $4a_1 \vdots a_0$, i.e. $a_1 = a_2 = \lambda a_0/4$, $a_3 = a_4 = (\lambda-2)a_0/2$, $d_1 = \lambda a_0/2$, $d_2 = (\lambda-2)a_0$,
- or $3a_1 \vdots a_0$, i.e. $a_1 = a_2 = \lambda a_0/3$, $a_3 = a_4 = (2\lambda/3-1)a_0$, $d_1 = 2\lambda a_0/3$, $d_2 = (4\lambda/3-2)a_0$.

If $a_0 = 3$ in the second case, then either the condition of Theorem QS (2), $(i,j) = (1,2)$, or Theorem WF is violated.

If $a_0 = 1$ in the first case, then by Theorem QS (2), $(i,j) = (1,2)$, $\lambda \in \{4, 8, 12\}$, i.e. $a_4 \le 5$.

If $a_0 = 2$ in the first case, then by Theorem QS (2), $(i,j) = (1,2)$, $\lambda = 6$, i.e. $a_4 \le 4$.

If $a_0 = 4$ in the first case, then the condition of Theorem QS (2), $(i,j) = (1,2)$ is violated.

Subcases $(13) = (23)$, $(4.1) - (4.9)$*:* The same analysis as in the previous subcases gives:

$$a_4 < 40.$$

Note that in Subcase (4.1) it may happen that $a_3 = a_4 = 2a_1 - a_0$, $d_1 = 2a_1$, $d_2 = 4a_1 - 2a_0$. Then by Theorem QS (2), $(i,j) = (1,2)$, either $a_1 = 3a_0$ or $a_1 = 2a_0$ or $a_1 = 3a_0/2$ or $a_1 = a_0$. In this case, $a_4 \le 5$ by Theorem WF and Lemma 2.

Case $(a_2 = a_3)$: *Subcase* (01)*:* $d_1 = a_0 + a_4$, $d_2 = a_1 + a_4$, $a_4 < 2a_2$ by (1). By Theorem WF,

- either $a_0 + a_4 \vdots a_2$, i.e. $a_4 = 2a_2 - a_0$ by (1),
- or $a_1 + a_4 \vdots a_2$, i.e. $a_4 = 2a_2 - a_1$ by (1).

If $a_4 = 2a_2 - a_1$, then $d_1 = 2a_2 + a_0 - a_1$, $d_2 = 2a_2$. By Theorem QS (2), $(i,j) = (2,3)$, $a_2 = a_3 = 2a_1 - a_0$. Then $d_1 = 3a_1 - a_0$, $d_2 = 4a_1 - 2a_0$ and by Theorem QS (1), $i = 0$,

- either $3a_1 \vdots a_0$, i.e. $a_1 = \lambda a_0/3$, $a_2 = a_3 = (2\lambda/3-1)a_0$, $a_4 = (\lambda-2)a_0$, $d_1 = (\lambda-1)a_0$, $d_2 = (4\lambda/3-2)a_0$,
- or $4a_1 \vdots a_0$, i.e. $a_1 = \lambda a_0/4$, $a_2 = a_3 = (\lambda/2-1)a_0$, $a_4 = (3\lambda/4-2)a_0$, $d_1 = (3\lambda/4-1)a_0$, $d_2 = (\lambda-2)a_0$.

In the second item $a_0 = 1$ by Theorem WF.

Hence

- either $a_0 = 3$, $a_1 = t$, $a_2 = a_3 = 2t - 3$, $a_4 = 3t - 6$, $d_1 = 3t - 3$, $d_2 = 4t - 6$, $t \not\equiv 0 \pmod 3$,
- or $a_0 = 1$, $a_1 = t$, $a_2 = a_3 = 2t - 1$, $a_4 = 3t - 2$, $d_1 = 3t - 1$, $d_2 = 4t - 2$.

These solutions appear in Table 1 (No. 17 and No. 26).

If $a_4 = 2a_2 - a_0$, then $d_1 = 2a_2$, $d_2 = 2a_2 + a_1 - a_0$. By Theorem QS (2), $(i,j) = (2,3)$, $a_1 = 2a_0$, and so $d_1 = 2a_2$, $d_2 = 2a_2 + a_0$. By Theorem QS (1), $i = 0$, $2a_2 \vdots a_0$, i.e.

$$a_1 = 2a_0,\ a_2 = a_3 = \lambda a_0/2,\ a_4 = (\lambda - 1)a_0,\ d_1 = \lambda a_0,\ d_2 = (\lambda + 1)a_0.$$

Hence

- either $a_0 = 2$, $a_1 = 4$, $a_2 = a_3 = 2t + 1$, $a_4 = 4t$, $d_1 = 4t + 2$, $d_2 = 4t + 4$,
- or $a_0 = 1$, $a_1 = 2$, $a_2 = a_3 = t$, $a_4 = 2t - 1$, $d_1 = 2t$, $d_2 = 2t + 1$, where t is odd.

These solutions appear in Table 1 (No. 16 and No. 24).

Subcase (02) = (03)*:* $d_1 = a_0 + a_4$, $d_2 = a_3 + a_4$, $a_4 < a_1 + a_3$ by (1). By Theorem WF,

- either $a_4 \vdots a_3$, i.e. $a_4 = a_3 = a_2$ by (1),
- or $a_0 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_0$ by (1).

By Lemma 3, we may assume that $a_4 = 2a_3 - a_0$, and so $d_1 = 2a_3$, $d_2 = 3a_3 - a_0$, $a_3 < a_0 + a_1$. By Theorem QS (1), $i = 1$,

- either $2a_3 \vdots a_1$, i.e. $a_3 = 3a_1/2$ by (1),
- or $3a_3 - a_0 \vdots a_1$, i.e. $a_3 = a_1 + a_0/3$ or $a_3 = 4a_1/3 + a_0/3$ by (1),
- or $2a_3 - a_0 \vdots a_1$, i.e. $a_3 = a_1 + a_0/2$ by (1).

If $a_3 = 3a_1/2$ or $a_3 = 4a_1/3 + a_0/3$, then $a_4 < 90$ by Theorem QS (1), $i = 0$, and Lemma 2.

If $a_3 = a_1 + a_0/3$, then $d_1 = 2a_1 + 2a_0/3$, $d_2 = 3a_1$. By Theorem QS (1), $i = 0$, $6a_1 \vdots a_0$, i.e.

$$a_1 = \frac{\lambda a_0}{6},\ a_2 = a_3 = \frac{\lambda + 2}{6}a_0,\ a_4 = \frac{\lambda - 1}{3}a_0,\ d_1 = \frac{\lambda + 2}{3}a_0,\ d_2 = \frac{\lambda a_0}{2}.$$

Hence

- either $a_0 = 6$, $a_1 = 2t + 1$, $a_2 = a_3 = 2t + 3$, $a_4 = 4t$, $d_1 = 4t + 6$, $d_2 = 6t + 3$, where $t \equiv 1 \pmod 3$,
- or $a_0 = 3$, $a_1 = t$, $a_2 = a_3 = t + 1$, $a_4 = 2t - 1$, $d_1 = 2t + 2$, $d_2 = 3t$, where $t \not\equiv -1 \pmod 3$.

These solutions appear in Table 1 (No. 28 and No. 40).

If $a_3 = a_1 + a_0/2$, then $a_4 = 2a_1$, $d_1 = 2a_1 + a_0$, $d_2 = 3a_1 + a_0/2$. By Theorem WF, $a_0 = a_1$. This situation was considered in a previous case.

Subcase (12) = (13)*:* $d_1 = a_1 + a_4$, $d_2 = a_3 + a_4$, $a_4 < a_0 + a_3$ by (1). By Theorem WF,

- either $a_4 \vdots a_3$, i.e. $a_4 = a_3 = a_2$ by (1),
- or $a_1 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_1$ by (1).

By Lemma 3, we may assume that $a_4 = 2a_3 - a_1$, and so $d_1 = 2a_3$, $d_2 = 3a_3 - a_1$, $a_3 < a_0 + a_1$. By Theorem QS (1), $i = 1$,

- either $2a_3 \vdots a_1$, i.e. $a_3 = 3a_1/2$ by (1),
- or $3a_3 \vdots a_1$, i.e. $a_3 = 4a_1/3$ or $a_3 = 5a_1/3$ by (1),
- or $3a_3 - a_0 \vdots a_1$, i.e. $a_3 = a_1 + a_0/3$ or $a_3 = 4a_1/3 + a_0/3$ by (1).

In each case, except for $a_3 = a_1 + a_0/3$, $a_4 < 96$ by Theorem QS (1), $i = 0$, and Lemma 2.

If $a_3 = a_1 + a_0/3$, then $d_1 = 2a_1 + 2a_0/3$, $d_2 = 2a_1 + a_0$. By Theorem QS (1), $i = 0$, $6a_1 \vdots a_0$, i.e.

$$a_1 = \frac{\lambda a_0}{6},\ a_2 = a_3 = \frac{\lambda+2}{6}a_0,\ a_4 = \frac{\lambda+4}{6}a_0,\ d_1 = \frac{\lambda+2}{3}a_0,\ d_2 = \frac{\lambda+3}{3}a_0.$$

Hence

- either $a_0 = 6$, $a_1 = 2t - 1$, $a_2 = a_3 = 2t + 1$, $a_4 = 2t + 3$, $d_1 = 4t + 2$, $d_2 = 4t + 4$, where $t \equiv -1 \pmod 3$,
- or $a_0 = 3$, $a_1 = t$, $a_2 = a_3 = t + 1$, $a_4 = t + 2$, $d_1 = 2t + 2$, $d_2 = 2t + 3$, where $3 \mid t$.

These solutions appear in Table 1 (No. 27 and No. 39).

Subcases (23), (4.1) – (4.9)*:* The same analysis gives:

$$a_4 < 30.$$

Case ($a_3 = a_4$): *Subcases* (01), (02), (12)*:* By Theorem WF, $d_1 \vdots a_4$ or $d_2 \vdots a_4$. Hence $a_2 = a_3 = a_4$, and we apply Lemma 3.

Subcase (03) = (4.1) = (4.4)*:* $d_1 = a_0 + a_4$, $d_2 = 2a_4$, $a_4 < a_1 + a_2$. By Theorem QS (1), $i = 2$,

- either $2a_4 \vdots a_2$, i.e. $a_4 = 3a_2/2$ by (1),
- or $a_0 + a_4 \vdots a_2$, i.e. $a_4 = 2a_2 - a_0$ by (1),
- or $a_0 + a_4 = a_e + \mu a_2$, $2a_4 = a_f + \nu a_2$ for some $e \neq f$, i.e. $(a_2 = 2a_1 - 3a_0,\ a_4 = 3a_1 - 4a_0)$ or $(a_1 = 3a_0/2,\ a_4 = a_2 + a_0/2)$.

If $a_4 = a_3 = 3a_2/2$, then $d_1 = a_0 + a_4 \vdots a_2/2$ by Theorem WF, i.e. $a_2 = 2a_0$, $a_4 = a_3 = 3a_0 \leq 3$ by Lemma 2.

If $a_4 = 2a_2 - a_0$, then $d_1 = 2a_2$, $d_2 = 4a_2 - 2a_0$, $a_2 < a_0 + a_1$. By Theorem QS (1), $i = 1$,

- either $2a_2 \vdots a_1$, i.e. $a_2 = 3a_1/2$ by (1),
- or $4a_2 - 2a_0 \vdots a_1$, i.e. $a_2 = \lambda a_1/4 + a_0/2$, $\lambda \in \{3, 4, 5\}$ by (1).

If $a_2 = 3a_1/2$, then $d_1 = 3a_1$, $d_2 = 6a_1 - 2a_0$, $a_1 < 2a_0$. By Theorem QS (1), $i = 0$, either $6a_1 \vdots a_0$ or $5a_1 \vdots a_0$ or $9a_1 \vdots a_0$. Then $a_4 < 45$ by Lemma 2.

If $a_2 = 5a_1/4 + a_0/2$, then $d_1 = 5a_1/2 + a_0$, $d_2 = 5a_1$, $a_1 < 2a_0$. By Theorem QS (1), $i = 0$, either $15a_1 \vdots a_0$ or $4a_1 \vdots a_0$. Then $a_4 < 60$ by Lemma 2.

If $a_2 = a_1 + a_0/2$, then $a_3 = a_4 = 2a_1$. By Theorem WF, $d_1 = 2a_1 + a_0 \vdots a_1$, i.e. $a_0 = a_1$, which is a previously considered case.

If $a_2 = 3a_1/4 + a_0/2$, then $a_3 = a_4 = 3a_1/2$. By Theorem WF, $d_1 = 3a_1/2 + a_0 \vdots a_1/2$, i.e. $a_1 = 2a_0$, and so $a_4 = a_3 = 3a_0 \leq 3$ by Lemma 2.

If $a_2 = 2a_1 - 3a_0$, $a_4 = 3a_1 - 4a_0$, then $d_1 = 3(a_1 - a_0)$, $d_2 = 6a_1 - 8a_0$. By Theorem QS (1), $i = 1$, either $9a_0 \vdots a_1$ or $8a_0 \vdots a_1$. Then $a_4 < 27$ by Lemma 2.

If $a_1 = 3a_0/2$, $a_4 = a_2 + a_0/2$, then $d_1 = a_2 + 3a_0/2$, $d_2 = 2a_2 + a_0$. By Theorem QS (1), $i = 1$, $6a_2 \vdots a_1$, i.e.

$$a_1 = \frac{3a_0}{2},\ a_2 = \frac{\lambda a_0}{4},\ a_3 = a_4 = \frac{\lambda+2}{4}a_0,\ d_1 = \frac{\lambda+6}{4}a_0,\ d_2 = \frac{\lambda+2}{2}a_0.$$

Hence

- either $a_0 = 2$, $a_1 = 3$, $a_2 = 3t$, $a_3 = a_4 = 3t + 1$, $d_1 = 3t + 3$, $d_2 = 6t + 2$,
- or $a_0 = 4$, $a_1 = 6$, $a_2 = 6t - 3$, $a_3 = a_4 = 6t - 1$, $d_1 = 6t + 3$, $d_2 = 12t - 2$.

These solutions appear in Table 1 (No. 22 and No. 32).

Subcase (13) = (4.2) = (4.5)*:* $d_1 = a_1 + a_4$, $d_2 = 2a_4$, $a_4 < a_0 + a_2$. By Theorem QS (1), $i = 2$,

- either $2a_4 \vdots a_2$, i.e. $a_4 = 3a_2/2$ by (1),
- or $a_1 + a_4 \vdots a_2$, i.e. $a_4 = 2a_2 - a_1$ by (1).

If $a_4 = 3a_2/2$, then $d_1 = a_1 + 3a_2/2$, $d_2 = 3a_2$, $a_2 < 2a_0$. By Theorem QS (1), $i = 1$,

- either $3a_2 \vdots a_1$, i.e. $a_2 = 4a_1/3$ or $a_2 = 5a_1/3$ by (1),
- or $2a_2 \vdots a_1$, i.e. $a_2 = 3a_1/2$ by (1),
- or $3a_2 - a_0 \vdots a_1$, i.e. $a_2 = a_1 + a_0/3$ or $a_2 = 4a_1/3 + a_0/3$ by (1).

In each case, $a_4 < 30$ by Theorem QS (1), $i = 0$, and Lemma 2.

If $a_4 = 2a_2 - a_1$, then $d_1 = 2a_2$, $d_2 = 4a_2 - 2a_1$, $a_2 < a_0 + a_1$. By Theorem QS (1), $i = 1$,

- either $4a_2 \vdots a_1$, i.e. $a_2 = \lambda a_1/4$, $\lambda \in \{5, 6, 7\}$ by (1),
- or $3a_2 \vdots a_1$, i.e. $a_2 = \lambda a_1/3$, $\lambda \in \{4, 5\}$ by (1),
- or $4a_2 - a_0 \vdots a_1$, i.e. $a_2 = \lambda a_1/4 + a_0/4$, $\lambda \in \{4, 5, 6\}$ by (1).

In each case, except for the last item with $\lambda = 4$, $a_4 < 50$ by Theorem QS (1), $i = 0$, and Lemma 2.

If $a_2 = a_1 + a_0/4$, then $d_1 = 2a_1 + a_0/2$, $d_2 = 2a_1 + a_0$. By Theorem QS (1), $i = 0$, $4a_1 \vdots a_0$, i.e. $a_1 = \lambda a_0/4$, $a_2 = (\lambda + 1)a_0/4$, $a_3 = a_4 = (\lambda + 2)a_0/4$. By Theorem WF,

$$a_0 = 4,\ a_1 = \lambda,\ a_2 = \lambda + 1,\ a_3 = a_4 = \lambda + 2,\ d_1 = 2\lambda + 2,\ d_2 = 2\lambda + 4.$$

By Theorem QS (2), $(i, j) = (0, 2)$, $\lambda = 4t + 1$. This solution appears in Table 1 (No. 37).

Subcase (23) = (4.3) = (4.6)*:* $d_1 = a_2 + a_4$, $d_2 = 2a_4$, $a_4 < a_0 + a_1$. By Theorem QS (1), $i = 2$,

- either $2a_4 \vdots a_2$, i.e. $a_4 = 3a_2/2$ by (1),
- or $2a_4 - a_0 \vdots a_2$, i.e. $a_4 = a_2 + a_0/2$ by (1),
- or $2a_4 - a_1 \vdots a_2$, i.e. $a_4 = a_2 + a_1/2$ by (1).

If $a_4 = a_2 + a_0/2$, then $d_1 = 2a_2 + a_0/2$, $d_2 = 2a_2 + a_0$, $a_2 < a_1 + a_0/2$. By Theorem QS (1), $i = 1$,

- either $2a_2 + a_0/2 \vdots a_1$, i.e. $a_2 = 3a_1/2 - a_0/4$ by (1),
- or $2a_2 + a_0 \vdots a_1$, i.e. $a_2 = 3a_1/2 - a_0/2$ by (1),
- or $2a_2 + a_0/2 = a_e + \mu a_1$, $2a_2 + a_0 = a_f + \nu a_1$ for some $e \neq f$, i.e. $a_1 = 5a_0/4$, $a_2 = 3a_0/2$ by (1).

In each case, $a_4 < 16$ by Theorem QS (1), $i = 0$, and Lemma 2.

If $a_4 = a_2 + a_1/2$, then $d_1 = 2a_2 + a_1/2$, $d_2 = 2a_2 + a_1$, $a_2 < a_0 + a_1/2$. By Theorem QS (1), $i = 1$, $4a_2 \vdots a_1$, i.e. $a_2 = \lambda a_1/4$, $\lambda < 6$, and so $a_4 = a_3 = (\lambda + 2)a_1/4 < 8$ by Lemma 2.

If $a_4 = 3a_2/2$, then $d_1 = 5a_2/2$, $d_2 = 3a_2$, $a_2 < 2(a_0 + a_1)/3$. By Theorem QS (1), $i = 1$, $5a_2 \vdots a_1$, i.e. $a_2 = 6a_1/5$. Then $a_4 = a_3 = 9a_1/5 \leq 9$ by Lemma 2.

Subcase (4.7) = (4.8) = (4.9)*:* $d_1 = d_2 = 2a_4$, $a_4 < (a_0 + a_1 + a_2)/2$. By Theorem QS (1), $i = 2$, $2a_4 \vdots a_2$, i.e. $a_2 = a_3 = a_4$ and we apply Lemma 3. *QED*

Lemma 5. *The* Main Theorem *holds in cases* (4.1) − (4.9) *of Lemma* 1.

Proof: By Lemma 4, we may assume that

$$1 \le a_0 \lneqq a_1 \lneqq a_2 \lneqq a_3 \lneqq a_4. \tag{3}$$

Case (4.1)*:* $d_1 = a_0 + a_3$, $d_2 = 2a_4$, $a_4 < a_1 + a_2$ by (1). By Theorem QS (1), $i = 3$,

- either $2a_4 \vdots a_3$, i.e. $a_4 = 3a_3/2$ by (1),
- or $2a_4 - a_2 \vdots a_3$, i.e. $a_4 = a_3 + a_2/2$ by (1),
- or $2a_4 - a_1 \vdots a_3$, i.e. $a_4 = a_3 + a_1/2$ by (1).

If $a_4 = 3a_3/2$, then $d_1 = a_0 + a_3$, $d_2 = 3a_3$, $a_3 < 2(a_1 + a_2)/3$. By Theorem QS (1), $i = 2$,

- either $3a_3 \vdots a_2$, i.e. $a_2 = a_3$ by (1),
- or $a_0 + a_3 \vdots a_2$, i.e. $a_3 = 2a_2 - a_0$ by (1),
- or $a_0 + a_3 = a_e + \mu a_2$, $3a_3 = a_f + \nu a_2$ for some $e \ne f$, i.e. $a_1 = 4a_0/3$, $a_3 = a_2 + a_0/3$.

In the last case, $d_1 = a_2 + 4a_0/3$, $d_2 = 3a_2 + a_0$, $a_2 < 5a_0/3$. By Theorem WF, $9a_2 \vdots a_0$, i.e. $a_2 = \lambda a_0/9$, $\lambda < 15$, $a_3 = (\lambda + 3)a_0/9$, and so $a_4 = (\lambda + 3)a_0/6 < 54$ by Lemma 2.

If $a_3 = 2a_2 - a_0$, then $d_1 = 2a_2$, $d_2 = 6a_2 - 3a_0$, $a_2 < a_1/2 + 3a_0/4$. By Theorem QS (1), $i = 1$,

- either $2a_2 \vdots a_1$, i.e. $a_1 = a_2$ by (1),
- or $6a_2 - 3a_0 \vdots a_1$, i.e. $a_2 = 2a_1/3 + a_0/2$ by (1).

The first case contradicts to (3). Hence $a_2 = 2a_1/3 + a_0/2$, and so $d_1 = 4a_1/3 + a_0$, $d_2 = 4a_1$, $a_3 = 4a_1/3$, $a_4 = 2a_1$. By Theorem WF, $a_0 \vdots \frac{a_1}{3}$. Then $a_4 \le 12$ by Lemma 2.

If $a_4 = a_3 + a_2/2$, then $d_1 = a_0 + a_3$, $d_2 = 2a_3 + a_2$, $a_3 < a_1 + a_2/2$. By Theorem QS (1), $i = 2$,

- either $2a_3 \vdots a_2$, i.e. $a_2 = a_3$ by (1),
- or $a_0 + a_3 \vdots a_2$, i.e. $a_3 = 2a_2 - a_0$ by (1),
- or $a_0 + a_3 = a_e + \mu a_2$, $2a_3 = a_f + \nu a_2$ for some $e \ne f$, i.e. $a_1 = 3a_0/2$, $a_3 = a_2 + a_0/2$.

In the second case $a_4 < 320$, while in the third case $a_4 < 51$ by Theorem QS, Theorem WF and Lemma 2.

If $a_4 = a_3 + a_1/2$, then $d_1 = a_0 + a_3$, $d_2 = 2a_3 + a_1$, $a_3 < a_2 + a_1/2$. By Theorem QS (1), $i = 2$,

- either $2a_3 + a_1 \vdots a_2$, i.e. $a_3 = 3a_2/2 - a_1/2$ by (1),
- or $a_0 + a_3 \vdots a_2$, i.e. $a_3 = 2a_2 - a_0$ by (1),
- or $a_0 + a_3 = a_e + \mu a_2$, $2a_3 + a_1 = a_f + \nu a_2$ for some $e \ne f$, i.e. $(a_2 = 3(a_1 - a_0)$, $a_3 = 4(a_1 - a_0))$ or $(a_2 = 2a_1 - a_0$, $a_3 = 3a_1 - 2a_0)$.

In the second case $a_4 < 5$, while in the third item $a_4 < 35$ by Theorem QS, Theorem WF and Lemma 2.

If $a_3 = 3a_2/2 - a_1/2$, then $d_1 = 3a_2/2 - a_1/2 + a_0$, $d_2 = 3a_2$, $a_2 < 2a_1$. By Theorem QS (1), $i = 1$,

- either $3a_2 \vdots a_1$, i.e. $a_2 = 5a_1/3$ or $a_2 = 4a_1/3$ by (1),

- or $3a_2 + 2a_0 \vdots a_1$, i.e. $a_2 = \lambda a_1/3 - 2a_0/3$, $\lambda \in \{4,5,6,7\}$ by (1),
- or $3a_2/2 - a_1/2 + a_0 = a_e + \mu a_1$, $3a_2 = a_f + \nu a_1$ for some $e \neq f$.

In the first item $a_4 < 16$, while in the third item $a_4 < 64$ by Lemma 2 by the same analysis as in the previous cases.

If $a_2 = 7a_1/3 - 2a_0/3$, then $d_1 = 3a_1$, $d_2 = 7a_1 - 2a_0$, $a_1 < 2a_0$. Then $a_4 < 168$ by Theorem QS (1), $i = 0$, and Lemma 2.

By Theorem QS (2), $(i,j) = (1,2)$, $d_1 = 3a_2/2 - a_1/2 + a_0 = a_e + \alpha a_1 + \beta a_2$, $\alpha, \beta \geq 0$, for some e. Together with (1), this implies that

- either $d_1 = a_1 + a_2$, i.e. $a_2 = 3a_1 - 2a_0$,
- or $d_1 = a_0 + 2a_1$, i.e. $a_2 = 5a_1/3$,
- or $d_1 = 2a_1$, i.e. $a_2 = 5a_1/3 - 2a_0/3$,
- or $d_1 = 3a_1$, i.e. $a_2 = 7a_1/3 - 2a_0/3$.

If $a_2 = 3a_1 - 2a_0 = \lambda a_1/3 - 2a_0/3$, then $a_4 < 18$ by Lemma 2. Hence it remains to consider the following case:

$$a_2 = \frac{5a_1}{3} - \frac{2a_0}{3},\ a_3 = 2a_1 - a_0,\ a_4 = \frac{5a_1}{2} - a_0,\ d_1 = 2a_1,\ d_2 = 5a_1 - 2a_0.$$

By Theorem QS (1), $i = 0$, either $4a_1 \vdots a_0$ or $10a_1 \vdots a_0$. Since $5 \nmid a_0$ by Theorem WF, $4a_1 \vdots a_0$. Hence

$$a_1 = \frac{\lambda a_0}{4},\ a_2 = \left(\frac{5\lambda}{12} - \frac{2}{3}\right) a_0,\ a_3 = \left(\frac{\lambda}{2} - 1\right) a_0,\ a_4 = \left(\frac{5\lambda}{8} - 1\right) a_0.$$

By Theorem WF,

$$a_0 = 1,\ a_1 = 2t,\ a_2 = \frac{2}{3}(5t-1),\ a_3 = 4t - 1,\ a_4 = 5t - 1,\ d_1 = 4t,\ d_2 = 10t - 2.$$

This violates the condition of Theorem QS (2), $(i,j) = (2,4)$.

Case (4.2)*:* $d_1 = a_1 + a_3$, $d_2 = 2a_4$, $a_4 < a_0 + a_2$ by (1). By Theorem QS (1), $i = 3$,

- either $2a_4 \vdots a_3$, i.e. $a_4 = 3a_3/2$ by (1),
- or $2a_4 - a_0 \vdots a_3$, i.e. $a_4 = a_3 + a_0/2$ by (1),
- or $2a_4 - a_2 \vdots a_3$, i.e. $a_4 = a_3 + a_2/2$ by (1).

In the first case $a_4 < 23$, while in the third case $a_4 < 39$ by Theorem QS and Lemma 2.

If $a_4 = a_3 + a_0/2$, then $d_1 = a_1 + a_3$, $d_2 = 2a_3 + a_0$, $a_3 < a_2 + a_0/2$. By Theorem QS (1), $i = 2$,

- either $a_1 + a_3 \vdots a_2$, i.e. $a_3 = 2a_2 - a_1$ by (1),
- or $a_0 + 2a_3 \vdots a_2$, i.e. $a_3 = 3a_2/2 - a_0/2$ by (1).

In the first case $a_4 < 120$, while in the second case $a_4 < 36$ by Theorem QS, Theorem WF and Lemma 2.

Cases (4.3) – (4.9)*:* By the same analysis, $a_4 < 75$. *QED*

Lemma 6. *The* Main Theorem *holds in cases* (03), (13) *and* (23) *of Lemma* 1.

Proof: We assume (3) and consider each case separately.

Case (03)*:* $d_1 = a_0 + a_4$, $d_2 = a_3 + a_4$, $a_4 < a_1 + a_2$ by (1). By Theorem QS (1), $i = 3$,

- either $a_4 \vdots a_3$, i.e. $a_4 = a_3$ by (1),
- or $a_0 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_0$ by (1),
- or $a_0 + a_4 - a_1 \vdots a_3$, i.e. $a_4 = a_3 + a_1 - a_0$ by (1),
- or $a_0 + a_4 - a_2 \vdots a_3$, i.e. $a_4 = a_3 + a_2 - a_0$ by (1).

If $a_4 = 2a_3 - a_0$, then $a_3 = a_2 + a_0/3$ by Theorem QS (1), $i = 2$. Then $d_1 = 2a_2 + 2a_0/3$, $d_2 = 3a_2$, $a_2 < a_1 + a_0/3$. By Theorem QS (1), $i = 1$,

- either $d_1 \vdots a_1$, i.e. $a_2 = 3a_1/2 - a_0/3$ by (1),
- or $d_1 - a_0 \vdots a_1$, i.e. $a_2 = a_1 + a_0/6$ by (1).

In the first case, $a_4 < 54$ by Theorem QS and Lemma 2.

In the second case, $a_4 = 2a_1$, $d_1 = 2a_1 + a_0$, $d_2 = 3a_1 + a_0/2$. By Theorem WF, $a_1 = a_0$, which contradicts to (3).

If $a_4 = a_3 + a_1 - a_0$, then $d_1 = a_1 + a_3$, $d_2 = 2a_3 + a_1 - a_0$, $a_3 < a_0 + a_2$. By Theorem QS (1), $i = 2$,

- either $a_1 + a_3 \vdots a_2$, i.e. $a_3 = 2a_2 - a_1$ by (1),
- or $2a_3 + a_1 - a_0 \vdots a_2$, i.e. $a_3 = (3a_2 + a_0 - a_1)/2$ by (1).

If $a_3 = 2a_2 - a_1$, then $d_1 = 2a_2$, $d_2 = 4a_2 - a_1 - a_0$, $a_2 < a_0 + a_1$. By Theorem QS (1), $i = 1$,

- either $2a_2 \vdots a_1$, i.e. $a_2 = 3a_1/2$ by (1),
- or $4a_2 - a_0 \vdots a_1$, i.e. $a_2 = \lambda a_1/4 + a_0/4$, $\lambda \in \{4, 5, 6\}$ by (1),
- or $4a_2 - 2a_0 \vdots a_1$, i.e. $a_2 = \lambda a_1/4 + a_0/2$, $\lambda \in \{3, 4, 5\}$ by (1),
- or $3a_2 - a_0 \vdots a_1$, i.e. $a_2 = (\lambda a_1 + a_0)/3$, $\lambda \in \{3, 4\}$ by (1).

Theorem QS (1), $i = 0$, and Lemma 2 give bound $a_4 < 195$ in all cases except for the following two:

- $a_2 = a_1 + a_0/4$,
- $a_2 = a_1 + a_0/3$.

In these cases, by Theorem QS and Theorem WF,

- either $a_0 = 8$, $a_1 = 4t + 1$, $a_2 = 4t + 3$, $a_3 = 4t + 5$, $a_4 = 8t - 2$, $d_1 = 8t + 6$, $d_2 = 12t + 3$,
- or $a_0 = 9$, $a_1 = t$, $a_2 = t + 3$, $a_3 = t + 6$, $a_4 = 2t - 3$, $d_1 = 2t + 6$, $d_2 = 3t + 3$, where $t \equiv -1 \pmod 3$.

These solutions appear in Table 1 (No. 43 and No. 45).

If $a_3 = (3a_2 + a_0 - a_1)/2$, then $d_1 = (3a_2 + a_0 + a_1)/2$, $d_2 = 3a_2$, $a_2 < a_0 + a_1$. By Theorem QS (1), $i = 1$,

- either $3a_2 \vdots a_1$, i.e. $a_2 = 4a_1/3$ or $a_2 = 5a_1/3$ by (1),
- or $3a_2 + a_0 \vdots a_1$, i.e. $a_2 = \lambda a_1/3 - a_0/3$, $\lambda \in \{4, 5, 6\}$ by (1),
- or $3a_2 - a_0 \vdots a_1$, i.e. $a_2 = \lambda a_1/3 + a_0/3$, $\lambda \in \{3, 4\}$ by (1),
- or $2a_2 \vdots a_1$, i.e. $a_2 = 3a_1/2$ by (1).

In each case, $a_4 < 220$ by Theorem WF, Theorem QS and Lemma 2.

If $a_4 = a_3 + a_2 - a_0$, then $d_1 = a_2 + a_3$, $d_2 = 2a_3 + a_2 - a_0$, $a_3 < a_0 + a_1$. By Theorem QS (1), $i = 2$, $2a_3 - a_0 \vdots a_2$, i.e. $a_3 = a_2 + a_0/2$ by (1). Then $d_1 = 2a_2 + a_0/2$, $d_2 = 3a_2$, $a_2 < a_1 + a_0/2$. By Theorem QS (1), $i = 1$,

- either $3a_2 \vdots a_1$, i.e. $a_2 = 4a_1/3$ by (1),
- or $4a_2 + a_0 \vdots a_1$, i.e. $a_2 = \lambda a_1/4 - a_0/4$, $\lambda \in \{5, 6\}$ by (1),
- or $2a_2 - a_0/2 \vdots a_1$, i.e. $a_2 = a_1 + a_0/4$ by (1).

In the first two items, $a_4 < 286$ by Theorem QS and Lemma 2. In the third item, $a_4 = 2a_1$, $d_1 = 2a_1 + a_0$, $d_2 = 3a_1 + 3a_0/4$. By Theorem WF, $3a_0 \vdots a_1$, i.e. $a_1 = 3a_0$ or $a_1 = 3a_0/2$. In each case, $a_4 < 25$ by Lemma 2.

Case (13)*:* $d_1 = a_1 + a_4$, $d_2 = a_3 + a_4$, $a_4 < a_0 + a_2$ by (1). By Theorem QS (1), $i = 3$,

- either $a_1 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_1$ by (1),
- or $a_1 + a_4 - a_2 \vdots a_3$, i.e. $a_4 = a_3 + a_2 - a_1$ by (1).

If $a_4 = 2a_3 - a_1$, then $a_3 = a_2 + a_1/3$ by Theorem QS (1), $i = 2$. Then $d_1 = 2a_2 + 2a_1/3$, $d_2 = 3a_2$, $a_2 < a_0 + a_1/3$. By Theorem QS (1), $i = 1$, $6a_2 \vdots a_1$, i.e. $a_2 = 7a_1/6$, $a_3 = 3a_1/2$, $a_4 = 2a_1 \le 12$ by Lemma 2.

If $a_4 = a_3 + a_2 - a_1$, then $d_1 = a_2 + a_3$, $d_2 = 2a_3 + a_2 - a_1$, $a_3 < a_0 + a_1$. By Theorem QS (1), $i = 2$,

- either $2a_3 - a_1 \vdots a_2$, i.e. $a_3 = a_2 + a_1/2$ by (1),
- or $2a_3 - a_1 - a_0 \vdots a_2$, i.e. $a_3 = (a_0 + a_1 + a_2)/2$ by (1),
- or $2a_3 - 2a_1 \vdots a_2$, i.e. $a_3 = a_1 + a_2/2$ by (1).

In each case, $a_4 < 288$ by Theorem WF, Theorem QS and Lemma 2.

Case (23)*:* $d_1 = a_2 + a_4$, $d_2 = a_3 + a_4$, $a_4 < a_0 + a_1$ by (1). By Theorem QS (1), $i = 3$, $a_2 + a_4 \vdots a_3$, i.e. $a_4 = 2a_3 - a_2$. Then $d_1 = 2a_3$, $d_2 = 3a_3 - a_2$, $a_3 < (a_0 + a_1 + a_2)/2$. By Theorem QS (1), $i = 2$,

- either $3a_3 \vdots a_2$, i.e. $a_3 = 4a_2/3$ by (1),
- or $3a_3 - a_0 \vdots a_2$, i.e. $a_3 = a_2 + a_0/3$ by (1),
- or $3a_3 - a_1 \vdots a_2$, i.e. $a_3 = a_2 + a_1/3$ by (1).

In each case, $a_4 < 42$ by Theorem WF, Theorem QS and Lemma 2. *QED*

Lemma 7. *Assume given integers* $1 \le a_0 < a_1 < a_2 < a_3 < a_4 < 2a_3$. *Then* $\forall\, i \ne j \in \{0, 1, 2\}$,

$$a_i + a_4 \in (a_3, a_4) \quad \textit{if and only if} \quad a_4 = 2a_3 - a_i,$$

$$a_i + a_4 - a_j \in (a_3, a_4) \quad \textit{if and only if} \quad a_4 = \begin{cases} 2a_3 + a_j - a_i & \textit{if } i > j, \\ a_3 + a_j - a_i & \textit{if } i < j. \end{cases}$$

Here $(a_3, a_4) = \{\alpha a_3 + \beta a_4 \mid \alpha, \beta \in \mathbb{Z},\ \alpha, \beta \ge 0\}$.

Proof: If $a_i + a_4 = \alpha a_3 + \beta a_4$, then $\beta = 0$ and then $\alpha a_3 = a_i + a_4 < 3a_3$. Hence $\alpha = 2$.

If $a_i + a_4 = a_j + \alpha a_3 + \beta a_4$, then $\beta = 0$. Then $\alpha a_3 = a_i - a_j + a_4 < 3a_3$ if $i > j$, and $\alpha a_3 < a_4 < 2a_3$ if $i < j$. Hence $\alpha = 2$ if $i > j$, and $\alpha = 1$ if $i < j$. *QED*

Corollary 2. *The condition of Theorem* QS (2), $(i, j) = (3, 4)$, *can be rewritten as follows in cases* (01), (02) *and* (12) *of Lemma* 1, *assuming* $a_i \ne a_j$ *for* $i \ne j$ *(and given the assumptions of Lemma* 1*):*

- *(01):*
  - *either* $a_4 = 2a_3 - a_0$,
  - *or* $a_4 = 2a_3 - a_1$,
  - *or* $a_2 = 2a_1 - a_0$, $a_4 = a_3 + a_1 - a_0$,
  - *or* $a_3 = a_2 + a_1 - 2a_0$, $a_4 = 2a_2 + a_1 - 3a_0$;
- *(02):*
  - *either* $a_4 = 2a_3 - a_0$,
  - *or* $a_4 = 2a_3 - a_2$,
  - *or* $a_3 = a_2 - a_0$, $a_4 = a_2 + a_1 - 2a_0$,
  - *or* $a_3 = a_2 + a_1 - 2a_0$, $a_4 = a_2 + 2a_1 - 3a_0$,
  - *or* $a_3 = 2a_2 - a_1 - a_0$, $a_4 = 3a_2 - a_1 - 2a_0$;
- *(12):*
  - *either* $a_4 = 2a_3 - a_1$,
  - *or* $a_4 = 2a_3 - a_2$,
  - *or* $a_2 = 2a_1 - a_0$, $a_4 = 2a_3 + a_0 - a_1$,
  - *or* $a_3 = 2a_2 - a_1 - a_0$, $a_4 = 3a_2 - 2a_1 - a_0$.

Proof: Let us consider Case (01). The other two cases are similar. The condition

$$d_1 - a_e,\ d_2 - a_f \in (a_3, a_4),\ \ \#\{e\} = \#\{f\} = 2,\ \{e, f\} = \{0, 1, 2\}$$

of Theorem QS is the same as:

- either $d_1 - a_2,\ d_2 - a_2 \in (a_3, a_4)$,
- or $d_1 - a_1,\ d_2 - a_2 \in (a_3, a_4)$,
- or $d_1 - a_2,\ d_2 - a_0 \in (a_3, a_4)$.

The first case can not occur, because $a_0 \neq a_1$. Hence by Lemma 7,

- either $a_4 = a_3 + a_1 - a_0 = a_3 + a_2 - a_1$,
- or $a_4 = a_3 + a_2 - a_0 = 2a_3 + a_0 - a_1$.

QED

4 Proof of the Main Theorem

By Lemma 5 and Lemma 6, it remains to consider cases (01), (02) and (12) of Lemma 1. Each of these cases splits into several subcases according to Corollary 2. Lemma 4 allows us to assume (3).

4.1 Case (01)

4.1.1 Subcase $a_4 = 2a_3 - a_0$

In this subcase, $d_1 = 2a_3$, $d_2 = 2a_3 + a_1 - a_0$, $a_3 < a_0 + a_2$. By Theorem QS (1), $i = 2$,

- either $2a_3 \vdots a_2$, i.e. $a_3 = 3a_2/2$ by (1),
- or $2a_3 + a_1 - a_0 \vdots a_2$, i.e. $a_3 = (3a_2 + a_0 - a_1)/2$ by (1),
- or $2a_3 - a_0 \vdots a_2$, i.e. $a_3 = a_2 + a_0/2$ by (1).

In each case, $a_4 < 198$ by Theorem QS, Theorem WF and Lemma 2.

4.1.2 Subcase $a_4 = 2a_3 - a_1$

In this subcase, $d_1 = 2a_3 + a_0 - a_1$, $d_2 = 2a_3$, $a_3 < a_1 + a_2$. By Theorem QS (1), $i = 2$,

- either $2a_3 \vdots a_2$, i.e. $a_3 = 3a_2/2$ by (1),
- or $2a_3 + a_0 - a_1 \vdots a_2$, i.e. $a_3 = (3a_2 + a_1 - a_0)/2$ or $a_3 = a_2 + (a_1 - a_0)/2$ by (1),
- or $a_3 + a_0 - a_1 \vdots a_2$, i.e. ($a_1 = 2a_0$, $a_3 = a_2 + a_0$) or ($a_1 = 3a_0/2$, $a_3 = a_2 + a_0/2$) or ($a_2 = 2a_1 - 3a_0$, $a_3 = 3a_1 - 4a_0$) by (1),
- or $2a_3 + a_0 - 2a_1 \vdots a_2$, i.e. $a_2 = 2(a_1 - a_0)$, $a_3 = 3a_1 - 5a_0/2$ by (1),
- or $2a_3 - a_1 \vdots a_2$, i.e. $a_3 = a_2 + a_1/2$ by (1).

If $a_3 = a_2 + a_1/2$, then $a_4 = 2a_2$, $d_1 = 2a_2 + a_0$, $d_2 = 2a_2 + a_1$. By Theorem WF, $a_1 = a_2$, which contradicts to (3).

If $a_2 = 2(a_1 - a_0)$, $a_3 = 3a_1 - 5a_0/2$, then $a_4 = 5(a_1 - a_0)$, $d_1 = a_0 + 5(a_1 - a_0)$, $d_2 = a_0 + 6(a_1 - a_0)$. Hence $a_0 \vdots (a_1 - a_0)$ by Theorem WF, i.e. $a_1 = \frac{\lambda+1}{\lambda} a_0$. Then $a_2 = \frac{2}{\lambda} a_0$, and so $a_4 = \frac{5}{\lambda} a_0 \leq 5$ by Lemma 2.

If $a_1 = 2a_0$, $a_3 = a_2 + a_0$, then $a_4 = 2a_2$, $d_1 = 2a_2 + a_0$, $d_2 = 2a_2 + 2a_0$. By Theorem WF, $2a_0 \vdots a_2$, i.e. $a_2 = 2a_0 = a_1$, which contradicts to (3).

If $a_2 = 2a_1 - 3a_0$, $a_3 = 3a_1 - 4a_0$, then $d_1 = 5a_1 - 7a_0$, $d_2 = 6a_1 - 8a_0$. By Theorem QS (1), $i = 1$, either $3a_0 \vdots a_1$ or $7a_0 \vdots a_1$ or $8a_0 \vdots a_1$, which implies that $a_4 < 40$ by Lemma 2.

If $a_1 = 3a_0/2$, $a_3 = a_2 + a_0/2$, then $d_1 = 2a_2 + a_0/2$, $d_2 = 2a_2 + a_0$. By Theorem WF, $2a_2 \vdots a_0/2$, i.e.

$$a_2 = \frac{\lambda a_0}{4},\ a_3 = \frac{\lambda+2}{4} a_0,\ a_4 = \frac{\lambda-1}{2} a_0,\ d_1 = \frac{\lambda+1}{2} a_0,\ d_2 = \frac{\lambda+2}{2} a_0.$$

Hence

- either $a_0 = 4$, $a_1 = 6$, $a_2 = t$, $a_3 = t + 2$, $a_4 = 2t - 2$, $d_1 = 2t + 2$, $d_2 = 2t + 4$, where $t \equiv -1 \pmod 3$ is odd,
- or $a_0 = 2$, $a_1 = 3$, $a_2 = t$, $a_3 = t + 1$, $a_4 = 2t - 1$, $d_1 = 2t + 1$, $d_2 = 2t + 2$, where $t \equiv 1 \pmod 3$.

These solutions appear in Table 1 (No. 23 and No. 31).

If $a_3 = 3a_2/2$, then $d_1 = 3a_2 + a_0 - a_1$, $d_2 = 3a_2$, $a_2 < 2a_1$. By Theorem QS (2), $(i, j) = (2, 3)$,

- either $a_1 - a_0 \vdots a_2/2$, i.e. $a_2 = 2(a_1 - a_0)$,
- or $a_0 \vdots a_2/2$, i.e. $a_2 = 2a_0$,
- or $2a_1 - a_0 \vdots a_2/2$, i.e. $a_2 = 2a_1 - a_0$ or $a_2 = (4a_1 - 2a_0)/3$.

If $a_2 = 2(a_1 - a_0)$, then $d_1 = 5(a_1 - a_0)$, $d_2 = 6(a_1 - a_0)$. By Theorem QS (1), $i = 1$, either $5a_0 \vdots a_1$ or $6a_0 \vdots a_1$. Then $a_4 < 30$ by Lemma 2.

If $a_2 = 2a_0$, then $d_1 = 7a_0 - a_1$, $d_2 = 6a_0$. By Theorem QS (1), $i = 1$, $N \cdot a_0 \vdots a_1$, where $N \in \{4, 5, 6, 7\}$. Then $a_4 < 35$ by Lemma 2.

If $a_2 = 2a_1 - a_0$, then $d_1 = 5a_1 - 2a_0$, $d_2 = 6a_1 - 3a_0$. By Theorem QS (1), $i = 1$, either $2a_0 \vdots a_1$ or $3a_0 \vdots a_1$. Then $a_4 < 15$ by Lemma 2.

If $a_2 = (4a_1 - 2a_0)/3$, then $d_1 = 3a_1 - a_0$, $d_2 = 4a_1 - 2a_0$. By Theorem QS (1), $i = 0$, either $3a_1 \vdots a_0$ or $8a_1 \vdots a_0$. By Theorem WF, $a_0 = 1$, i.e.

$$a_0 = 1,\ a_1 = 2t + 1,\ a_2 = \frac{8t+2}{3},\ a_3 = 4t + 1,\ a_4 = 6t + 1,\ d_1 = 6t + 2,\ d_2 = 8t + 2.$$

This solution appears in Table 1 (No. 20).

If $a_3 = (3a_2 + a_1 - a_0)/2$, then $d_1 = 3a_2$, $d_2 = 3a_2 + a_1 - a_0$, $a_2 < a_0 + a_1$. By Theorem QS (1), $i = 1$,

- either $3a_2 \vdots a_1$, i.e. $a_2 = 4a_1/3$ or $a_2 = 5a_1/3$ by (1),
- or $3a_2 - a_0 \vdots a_1$, i.e. $a_2 = (\lambda a_1 + a_0)/3$, $\lambda \in \{3, 4\}$ by (1),
- or $d_1 - a_3 \vdots a_1$, i.e. $a_2 = (\lambda a_1 - a_0)/3$, $\lambda \in \{4, 5, 6\}$ by (1),
- or $2a_2 \vdots a_1$, i.e. $a_2 = 3a_1/2$ by (1).

In each case, $a_4 < 308$ by Theorem WF, Theorem QS and Lemma 2.

If $a_3 = a_2 + (a_1 - a_0)/2$, then $d_1 = 2a_2$, $d_2 = 2a_2 + a_1 - a_0$. By Theorem QS (1), $i = 1$,

- either $2a_2 \vdots a_1$, i.e. $a_2 = \lambda a_1/2$, $a_3 = (\lambda+1)a_1/2 - a_0/2$, $a_4 = \lambda a_1 - a_0$, $d_1 = \lambda a_1$, $d_2 = (\lambda+1)a_1 - a_0$,
- or $2a_2 - a_0 \vdots a_1$, i.e. $a_2 = (\lambda a_1 + a_0)/2$, $a_3 = (\lambda + 1)a_1/2$, $a_4 = \lambda a_1$, $d_1 = \lambda a_1 + a_0$, $d_2 = (\lambda + 1)a_1$,
- or $2a_2 - a_3 \vdots a_1$, i.e. $a_2 = \lambda a_1 + (a_1 - a_0)/2$, $a_3 = (\lambda+1)a_1 - a_0$, $a_4 = (2\lambda+1)a_1 - 2a_0$, $d_1 = (2\lambda+1)a_1 - a_0$, $d_2 = (2\lambda + 2)a_1 - 2a_0$.

In the second case, $2a_0 \vdots a_1$ by Theorem WF. Hence

$$a_1 = 2a_0,\ a_2 = \left(\lambda + \frac{1}{2}\right) a_0,\ a_3 = (\lambda + 1)a_0,\ a_4 = 2\lambda a_0.$$

This violates conditions of Theorem WF.

In the first case, $\gcd(a_0, a_1) = 1$ if λ is even, and $\gcd(a_0, a_1) = 2$ if λ is odd. In the third case, $\gcd(a_0, a_1) = 1$ by Theorem WF. In these two cases, by Theorem QS (1), $i = 0$,

- either $\lambda a_1 \vdots a_0$ or $(\lambda + 1)a_1 \vdots a_0$ or $((\lambda/2 + 1) \vdots a_0$ and λ is even) in the first case;
- and either $(2\lambda + 1) \vdots a_0$ or $(\lambda + 1) \vdots a_0$ or $(2\lambda + 3) \vdots a_0$ in the third case.

If $\lambda a_1 \vdots a_0$ in the first case, then $d_2 - a_1 = (\lambda a_1/a_0 - 1)a_0 \in (a_0, a_2, a_4)$. Hence by Theorem QS (3), $(k, l, m) = (0, 2, 4)$, either $d_2 = 2a_3 \in (a_0, a_2, a_4)$ or $a_3 = d_2 - a_3 \in (a_0, a_2, a_4)$, i.e. $2a_3 \in (a_0, a_2, a_4)$. Hence $2a_3 \in (a_0, a_2)$. This implies that either $(\lambda + 1)a_1 \vdots a_0$ or $((\lambda/2 + 1) \vdots a_0$ and λ is even).

If $(2\lambda + 1) \vdots a_0$ in the third case, then $d_2 - a_1 = a_4 \in (a_0, a_2, a_4)$. Hence by Theorem QS (3), $(k, l, m) = (0, 2, 4)$, $d_2 = 2a_3 \in (a_0, a_2, a_4)$. Hence $2a_3 \in (a_0, a_2)$. This implies that either $(\lambda + 1) \vdots a_0$ or $(2\lambda + 3) \vdots a_0$.

Hence

- either $a_0 = 2b_0$, $a_1 = 2b_1$, $a_2 = (\nu b_0 - 1)b_1$, $a_3 = (\nu b_1 - 1)b_0$, $a_4 = 2(\nu b_0 b_1 - b_0 - b_1)$, $d_1 = 2b_1(\nu b_0 - 1)$, $d_2 = 2b_0(\nu b_1 - 1)$, where $\gcd(b_0, b_1) = 1$, b_0, b_1 are odd, ν is even;
- or $a_2 = (\nu a_0 - 1)a_1/2$, $a_3 = (\nu a_1 - 1)a_0/2$, $a_4 = \nu a_0 a_1 - a_0 - a_1$, $d_1 = a_1(\nu a_0 - 1)$, $d_2 = a_0(\nu a_1 - 1)$, where $\gcd(a_0, a_1) = 1$, ν, a_0, a_1 are odd;
- or $a_2 = (\nu a_0 - 1)a_1$, $a_3 = \nu a_0 a_1 - (a_0 + a_1)/2$, $a_4 = 2\nu a_0 a_1 - a_0 - 2a_1$, $d_1 = 2a_1(\nu a_0 - 1)$, $d_2 = 2\nu a_0 a_1 - a_0 - a_1$, where $\gcd(a_0, a_1) = 1$, a_0, a_1 are odd;

- or $a_2 = (\nu a_0 - 1)a_1 + (a_1 - a_0)/2$, $a_3 = a_0(\nu a_1 - 1)$, $a_4 = 2\nu a_0 a_1 - 2a_0 - a_1$, $d_1 = 2\nu a_0 a_1 - a_0 - a_1$, $d_2 = 2a_0(\nu a_1 - 1)$, where $\gcd(a_0, a_1) = 1$, a_0, a_1 are odd;
- or $a_2 = (\nu a_1 - 1)a_0/2 - a_1$, $a_3 = (\nu a_0 - 1)a_1/2 - a_0$, $a_4 = \nu a_0 a_1 - 2(a_0 + a_1)$, $d_1 = \nu a_0 a_1 - a_0 - 2a_1$, $d_2 = \nu a_0 a_1 - 2a_0 - a_1$, where $\gcd(a_0, a_1) = 1$, ν, a_0, a_1 are odd.

These solutions appear in Table 1 (Nos. 10-14).

4.1.3 Subcase $a_2 = 2a_1 - a_0$, $a_4 = a_3 + a_1 - a_0$

In this subcase, $d_1 = a_3 + a_1$, $d_2 = a_3 + 2a_1 - a_0 = a_2 + a_3$. By Theorem QS (1), $i = 2$,

- either $a_3 \vdots a_2$, i.e. $a_3 = (\mu - 1)(2a_1 - a_0)$,
- or $a_3 + a_1 \vdots a_2$, i.e. $a_3 = \mu(2a_1 - a_0) - a_1$,
- or $a_1 + a_3 - a_0 \vdots a_2$, i.e. $a_4 \vdots a_2$.

If $a_4 \vdots a_2$, then by Theorem WF, either $d_1 = a_0 + a_4 \vdots a_2$ or $d_2 = a_1 + a_4 \vdots a_2$, i.e. $a_1 = a_2$. This contradicts to (3).

By Theorem QS (1), $i = 0$,

- either $a_3 + a_1 \vdots a_0$, i.e. $a_3 = \lambda a_0 - a_1$,
- or $a_3 + 2a_1 \vdots a_0$, i.e. $a_3 = \lambda a_0 - 2a_1$,
- or $d_2 - a_2 \vdots a_0$, i.e. $a_3 \vdots a_0$,
- or $d_2 - a_3 \vdots a_0$, i.e. $2a_1 \vdots a_0$, and hence $a_1 \vdots a_0$.

Note that a_0 is odd and $\gcd(a_0, a_1) = 1$ by Theorem WF. Moreover, $a_3 \vdots a_0$ if and only if $a_1 \vdots a_0$ if and only if $a_0 = 1$ by Theorem WF.

If $a_3 = \lambda a_0 - 2a_1 = (\mu - 1)(2a_1 - a_0)$, then $\mu = \nu a_0$,

$$a_2 = 2a_1 - a_0,\ a_3 = (\nu a_0 - 1)(2a_1 - a_0),\ a_4 = \nu a_0(2a_1 - a_0) - a_1,\ d_1 = (\nu a_0 - 1)(2a_1 - a_0) + a_1,$$
$$d_2 = \nu a_0(2a_1 - a_0),\ \text{where } \gcd(\nu a_0 - 1, a_1 - a_0) = 1.$$

If $a_3 = \lambda a_0 - 2a_1 = \mu(2a_1 - a_0) - a_1$, then $\mu = (\nu a_0 - 1)/2$, ν is odd,

$$a_2 = 2a_1 - a_0,\ a_3 = \left(\frac{\nu a_0 - 1}{2}\right)(2a_1 - a_0) - a_1,\ a_4 = \left(\frac{\nu a_0 - 1}{2}\right)(2a_1 - a_0) - a_0,$$
$$d_1 = \left(\frac{\nu a_0 - 1}{2}\right)(2a_1 - a_0),\ d_2 = \left(\frac{\nu a_0 + 1}{2}\right)(2a_1 - a_0) - a_1,\ \text{where } \gcd\left(\frac{\nu a_0 - 3}{2}, a_1 - a_0\right) = 1.$$

If $a_3 = \lambda a_0 - a_1 = (\mu - 1)(2a_1 - a_0)$, then $\mu = (\nu a_0 + 1)/2$, ν is odd,

$$a_2 = 2a_1 - a_0,\ a_3 = \left(\frac{\nu a_0 - 1}{2}\right)(2a_1 - a_0),\ a_4 = \left(\frac{\nu a_0 - 1}{2}\right)(2a_1 - a_0) + a_1 - a_0,$$
$$d_1 = \left(\frac{\nu a_0 - 1}{2}\right)(2a_1 - a_0) + a_1,\ d_2 = \left(\frac{\nu a_0 + 1}{2}\right)(2a_1 - a_0),\ \text{where } \gcd\left(\frac{\nu a_0 - 1}{2}, a_1 - a_0\right) = 1.$$

If $a_3 = \lambda a_0 - a_1 = \mu(2a_1 - a_0) - a_1$, then $\mu = \nu a_0$,

$$a_2 = 2a_1 - a_0,\ a_3 = \nu a_0(2a_1 - a_0) - a_1,\ a_4 = \nu a_0(2a_1 - a_0) - a_0,\ d_1 = \nu a_0(2a_1 - a_0),$$
$$d_2 = (\nu a_0 + 1)(2a_1 - a_0) - a_1,\ \text{where } \gcd(\nu a_0 - 1, a_1 - a_0) = 1.$$

In all these cases a_0 is odd and $\gcd(a_0, a_1) = 1$.

These solutions appear in Table 1 (Nos. 6-9).

4.1.4 Subcase $a_3 = a_2 + a_1 - 2a_0$, $a_4 = 2a_2 + a_1 - 3a_0$

In this subcase, $d_1 = 2a_2 + a_1 - 2a_0$, $d_2 = 2a_2 + 2a_1 - 3a_0$. By Theorem QS (1), $i = 2$,

- either $2a_1 - 4a_0 \vdots a_2$, i.e. $a_2 = 2a_1 - 4a_0$,
- or $2a_1 - 3a_0 \vdots a_2$, i.e. $a_2 = 2a_1 - 3a_0$,
- or $a_1 - 3a_0 \vdots a_2$, i.e. $a_1 = 3a_0$.

In each case, $a_4 < 55$ by Theorem WF, Theorem QS and Lemma 2.

4.2 Case (02)

4.2.1 Subcase $a_4 = 2a_3 - a_0$

In this subcase, $d_1 = 2a_3$, $d_2 = 2a_3 + a_2 - a_0$, $a_3 < a_0 + a_1$. By Theorem QS (1), $i = 2$,

- either $2a_3 \vdots a_2$, i.e. $a_3 = 3a_2/2$ by (1),
- or $2a_3 - a_0 \vdots a_2$, i.e. $a_3 = a_2 + a_0/2$ by (1),
- or $2a_3 - a_1 \vdots a_2$, i.e. $a_3 = a_2 + a_1/2$ by (1).

In each case, $a_4 < 90$ by Theorem QS and Lemma 2.

4.2.2 Subcase $a_4 = 2a_3 - a_2$

In this subcase, $d_1 = 2a_3 + a_0 - a_2$, $d_2 = 2a_3$, $a_3 < a_1 + a_2$. By Theorem QS (1), $i = 2$,

- either $2a_3 \vdots a_2$, i.e. $a_3 = 3a_2/2$ by (1),
- or $2a_3 + a_0 \vdots a_2$, i.e. $a_3 = (3a_2 - a_0)/2$ or $a_3 = 2a_2 - a_0/2$ by (1),
- or $a_3 + a_0 \vdots a_2$, i.e. $a_3 = 2a_2 - a_0$ by (1),
- or $d_1 - a_1 \vdots a_2$, i.e. $a_3 = a_2 + (a_1 - a_0)/2$ or $a_3 = (3a_2 + a_1 - a_0)/2$ by (1).

If $a_3 = (3a_2 + a_1 - a_0)/2$, then $d_1 = 2a_2 + a_1$, $d_2 = 3a_2 + a_1 - a_0$, $a_2 < a_1 + a_0$. By Theorem QS (1), $i = 1$,

- either $2a_2 \vdots a_1$, i.e. $a_2 = 3a_1/2$ by (1),
- or $3a_2 - a_0 \vdots a_1$, i.e. $a_2 = a_1 + a_0/3$ or $a_2 = (4a_1 + a_0)/3$ by (1),
- or $2a_2 - a_0 \vdots a_1$, i.e. $a_2 = a_1 + a_0/2$ by (1).

If $a_2 = 3a_1/2$ or $a_2 = (4a_1 + a_0)/3$, then $a_4 < 154$ by Lemma 2 and Theorem QS.

If $a_2 = a_1 + a_0/3$, then $a_3 = 2a_1$, $a_4 = 3a_1 - a_0/3$, $d_1 = 3a_1 + 2a_0/3$, $d_2 = 4a_1$. By Theorem QS (2), $(i, j) = (1, 3)$, $2a_0 \vdots a_1$, i.e. $a_1 = 2a_0$, $a_3 = 4a_0$, and so $a_4 = 17a_0/3 \leq 17$ by Lemma 2.

If $a_2 = a_1 + a_0/2$, then $a_4 = 3a_1$, $d_1 = 3a_1 + a_0$, $d_2 = 4a_1 + a_0/2$. By Theorem WF, $a_0 \vdots a_1$, i.e. $a_0 = a_1$, which contradicts to (3).

If $a_3 = a_2 + (a_1 - a_0)/2$, then $d_1 = a_2 + a_1$, $d_2 = 2a_2 + a_1 - a_0$. By Theorem QS (1), $i = 1$,

- either $a_2 \vdots a_1$, i.e. $a_2 = \lambda a_1$, $a_3 = (\lambda + 1/2)a_1 - a_0/2$, $a_4 = (\lambda + 1)a_1 - a_0$, $d_1 = (\lambda + 1)a_1$, $d_2 = (2\lambda + 1)a_1 - a_0$,

- or $2a_2-a_0 \vdots a_1$, i.e. $a_2=(\lambda a_1+a_0)/2$, $a_3=(\lambda+1)a_1/2$, $a_4=(\lambda+2)a_1/2-a_0/2$, $d_1=(\lambda+2)a_1/2+a_0/2$, $d_2=(\lambda+1)a_1$,
- or $2a_2-2a_0 \vdots a_1$, i.e. $a_2=\lambda a_1/2+a_0$, $a_3=(\lambda+1)a_1/2+a_0/2$, $a_4=(\lambda+2)a_1/2$, $d_1=(\lambda+2)a_1/2+a_0$, $d_2=(\lambda+1)a_1+a_0$.

Note that in the first item, $\gcd(a_0,a_1)=1$, a_0, a_1 are odd. In the second item, $\gcd(a_0,a_1)\in\{1,2\}$ if λ is odd, and $\gcd(a_0,a_1)=2$ if λ is even.

In the third item, $a_1=2a_0$ by Theorem QS (2), $(i,j)=(1,4)$. Then a_0 should be even, which violates conditions of Theorem WF.

By Theorem QS (1), $i=0$,

- either $\lambda+1 \vdots a_0$ or $2\lambda+1 \vdots a_0$ or $\lambda \vdots a_0$ in the first item;
- and either $(\lambda+1)a_1 \vdots a_0$ or $\lambda a_1 \vdots a_0$ or ($(\lambda+2)a_1 \vdots a_0$ and $(\lambda+2)a_1/a_0$ is odd) in the second item.

In the first item, $\lambda \vdots a_0$ implies that either $\lambda+1 \vdots a_0$ or $2\lambda+1 \vdots a_0$ by Theorem QS (2), $(i,j)=(0,2)$. In the second item, $\lambda a_1 \vdots a_0$ implies that either $(\lambda+1)a_1 \vdots a_0$ or ($(\lambda+2)a_1 \vdots a_0$ and $(\lambda+2)a_1/a_0$ is odd) by Theorem QS (2), $(i,j)=(0,2)$.

Hence

- either $a_2=(\nu a_0-1)a_1$, $a_3=\nu a_0a_1-(a_0+a_1)/2$, $a_4=(\nu a_1-1)a_0$, $d_1=\nu a_0a_1$, $d_2=2\nu a_0a_1-a_0-a_1$, where $\gcd(a_0,a_1)=\gcd(\nu a_0-1,(a_1-a_0)/2)=1$, a_0, a_1 are odd;
- or $a_2=(\nu a_0-1)a_1/2$, $a_3=(\nu a_1-1)a_0/2$, $a_4=(\nu a_0+1)a_1/2-a_0$, $d_1=(\nu a_0+1)a_1/2$, $d_2=(\nu a_1-1)a_0$, where $\gcd(a_0,a_1)=\gcd((\nu a_0-1)/2,(a_1-a_0)/2)=1$, ν, a_0, a_1 are odd;
- or $a_0=2b_0$, $a_1=2b_1$, $a_2=(\nu b_0-1)b_1+b_0$, $a_3=\nu b_0b_1$, $a_4=(\nu b_0+1)b_1-b_0$, $d_1=(\nu b_0+1)b_1+b_0$, $d_2=2\nu b_0b_1$, where $\gcd(b_0,b_1)=\gcd(\nu,b_1-b_0)=1$, ν, b_0, b_1 are odd;
- or $a_2=\nu a_0a_1+(a_0-a_1)/2$, $a_3=\nu a_0a_1$, $a_4=\nu a_0a_1+(a_1-a_0)/2$, $d_1=\nu a_0a_1+(a_0+a_1)/2$, $d_2=2\nu a_0a_1$, where $\gcd(a_0,a_1)=\gcd(\nu,(a_1-a_0)/2)=1$, a_0, a_1 are odd;
- or $a_2=(\nu a_1+1)a_0/2-a_1$, $a_3=(\nu a_0-1)a_1/2$, $a_4=(\nu a_1-1)a_0/2$, $d_1=(\nu a_1+1)a_0/2$, $d_2=(\nu a_0-1)a_1$, where $\gcd(a_0,a_1)=\gcd((\nu a_0-1)/2,(a_1-a_0)/2)=1$, ν, a_0, a_1 are odd.

These solutions appear in Table 1 (Nos. 1-5).

If $a_3=3a_2/2$, then $a_4=2a_2$, $d_1=2a_2+a_0$, $d_2=3a_2$. Hence $a_0 \vdots a_2/2$ by Theorem WF, i.e. $a_2=2a_0$. Then $a_3=3a_0$, $a_4=4a_0\le 4$ by Lemma 2.

If $a_3=(3a_2-a_0)/2$, then $d_1=2a_2$, $d_2=3a_2-a_0$, $a_2<2a_1+a_0$. By Theorem QS (1), $i=1$,

- either $2a_2 \vdots a_1$, i.e. $a_2=\lambda a_1/2$, $\lambda\in\{3,4,5\}$ by (1),
- or $3a_2-a_0 \vdots a_1$, i.e. $a_2=(\lambda a_1+a_0)/3$, $3\le\lambda\le 7$ by (1),
- or $2a_2-a_0 \vdots a_1$, i.e. $a_2=(\lambda a_1+a_0)/2$, $\lambda\in\{2,3,4\}$ by (1),
- or $3a_2-2a_0 \vdots a_1$ and $2a_2-a_3 \vdots a_1$, i.e. $a_2=2a_1-a_0$ and $a_1=5a_0/\mu$, $1\le\mu\le 4$ by (1).

In the last item, $a_4<20$ by Lemma 2. In the second item, $4a_0 \vdots a_1$ by Theorem QS (2), $(i,j)=(1,3)$. Then $a_4<56$ by Lemma 2. In the third item, $a_4=\lambda a_1$, $d_1=\lambda a_1+a_0$, $d_2=(3\lambda a_1+a_0)/2$. Hence $a_0=a_1$ by Theorem WF, which violates (3).

If $a_2=5a_1/2$, then $a_4<450$ by Theorem QS and Lemma 2.

If $a_2 = 3a_1/2$, then $d_1 = 3a_1$, $d_2 = 9a_1/2 - a_0$. By Theorem QS (1), $i = 0$, either $9a_1 \vdots a_0$ or $7a_1 \vdots a_0$. Since $3 \nmid a_0$ by Theorem WF, $7a_1 \vdots a_0$, i.e.

$$a_1 = \frac{\lambda a_0}{7},\ a_2 = \frac{3\lambda a_0}{14},\ a_3 = \left(\frac{9\lambda}{28} - \frac{1}{2}\right) a_0,\ a_4 = \left(\frac{3\lambda}{7} - 1\right) a_0,\ d_1 = \frac{3\lambda a_0}{7},\ d_2 = \left(\frac{9\lambda}{14} - 1\right) a_0.$$

Hence

- either $a_0 = 1$, $a_1 = 2t$, $a_2 = 3t$, $a_3 = (9t-1)/2$, $a_4 = 6t - 1$, $d_1 = 6t$, $d_2 = 9t - 1$, where t is odd,
- or $a_0 = 7$, $a_1 = 2t$, $a_2 = 3t$, $a_3 = (9t-7)/2$, $a_4 = 6t - 7$, $d_1 = 6t$, $d_2 = 9t - 7$, where t is odd, $7 \nmid t$.

These solutions appear in Table 1 (No. 18 and No. 41).

If $a_2 = 2a_1$, then $d_1 = 4a_1$, $d_2 = 6a_1 - a_0$. By Theorem QS (1), $i = 0$, either $6a_1 \vdots a_0$ or $5a_1 \vdots a_0$ or $4a_1 \vdots a_0$. Since $\gcd(a_0, a_1) = 1$ by Theorem WF and a_0 is even, $a_0 \in \{2, 4, 6\}$, i.e.

- either $a_0 = 2$, $a_2 = 2a_1$, $a_3 = 3a_1 - 1$, $a_4 = 4a_1 - 2$, $d_1 = 4a_1$, $d_2 = 6a_1 - 2$,
- or $a_0 = 4$, $a_2 = 2a_1$, $a_3 = 3a_1 - 2$, $a_4 = 4a_1 - 4$, $d_1 = 4a_1$, $d_2 = 6a_1 - 4$,
- or $a_0 = 6$, $a_2 = 2a_1$, $a_3 = 3a_1 - 3$, $a_4 = 4a_1 - 6$, $d_1 = 4a_1$, $d_2 = 6a_1 - 6$.

In the first and the third items, either a_1 or a_3 is even, which violates conditions of Theorem WF. Hence

$$a_0 = 4,\ a_1 = 2t + 1,\ a_2 = 4t + 2,\ a_3 = 6t + 1,\ a_4 = 8t,\ d_1 = 8t + 4,\ d_2 = 12t + 2.$$

This solution appears in Table 1 (No. 36).

If $a_3 = 2a_2 - a_0$, then $d_1 = 3a_2 - a_0$, $d_2 = 4a_2 - 2a_0$, $a_2 < a_0 + a_1$. By Theorem QS (1), $i = 1$,

- either $3a_2 - a_0 \vdots a_1$, i.e. $a_2 = a_1 + a_0/3$ or $a_2 = (4a_1 + a_0)/3$ by (1),
- or $4a_2 - 2a_0 \vdots a_1$, i.e. $a_2 = \lambda a_1/4 + a_0/2$, $\lambda \in \{3, 4, 5\}$ by (1),
- or $3a_2 - 2a_0 \vdots a_1$, i.e. $a_2 = a_1 + 2a_0/3$ or $a_2 = 2(a_1 + a_0)/3$ by (1).

If $a_2 = (4a_1 + a_0)/3$ or $a_2 = 5a_1/4 + a_0/2$, then $a_4 < 420$ by Theorem QS and Lemma 2. If $a_2 = 3a_1/4 + a_0/2 > a_1$ or $a_2 = 2(a_1 + a_0)/3 > a_1$, then $a_1 < 2a_0$. Then $a_4 < 144$ by Theorem QS and Lemma 2.

If $a_2 = a_1 + a_0/3$, then $d_1 = 3a_1$, $d_2 = 4a_1 - 2a_0/3$. By Theorem QS (1), $i = 0$, either $12a_1 \vdots a_0$ or $9a_1 \vdots a_0$. Since $\gcd(a_0/3, a_1) = 1$ by Theorem WF, $a_0/3 \in \{1, 2, 3, 4\}$. Note that if $6 \mid a_0$, then the condition of Theorem QS (1), $i = 0$, is violated. Hence

- either $a_0 = 3$, $a_2 = a_1 + 1$, $a_3 = 2a_1 - 1$, $a_4 = 3a_1 - 3$, $d_1 = 3a_1$, $d_2 = 4a_1 - 2$, where $a_1 \equiv 1 \pmod 3$,
- or $a_0 = 9$, $a_2 = a_1 + 3$, $a_3 = 2a_1 - 3$, $a_4 = 3a_1 - 9$, $d_1 = 3a_1$, $d_2 = 4a_1 - 6$, where $a_1 \equiv -1 \pmod 3$.

These solutions appear in Table 1 (No. 29 and No. 44).

If $a_2 = a_1 + 2a_0/3$, then $a_4 = 3a_1$, $d_1 = 3a_1 + a_0$, $d_2 = 4a_1 + 2a_0/3$. By Theorem WF, $2a_0 \vdots a_1$, i.e. $a_1 = 2a_0$, $a_2 = 8a_0/3$, and so $a_4 = 6a_0 \leq 18$ by Lemma 2.

If $a_2 = a_1 + a_0/2$, then $d_1 = 3a_1 + a_0/2$, $d_2 = 4a_1$. By Theorem QS (1), $i = 0$, either $6a_1 \vdots a_0$ or $4a_1 \vdots a_0$. Since $\gcd(a_0/2, a_1) = 1$ by Theorem WF, $a_0/2 \in \{1, 2, 3\}$. Note that if $a_0/2$ is odd, then a condition of Theorem WF is violated. Hence

$$a_0 = 4,\ a_1 = 2t + 1,\ a_2 = 2t + 3,\ a_3 = 4t + 2,\ a_4 = 6t + 1,\ d_1 = 6t + 5,\ d_2 = 8t + 4.$$

This solution appears in Table 1 (No. 33).

If $a_3 = 2a_2 - a_0/2$, then $d_1 = 3a_2$, $d_2 = 4a_2 - a_0$, $a_2 < a_1 + a_0/2$. By Theorem QS (1), $i = 1$,

- either $3a_2 \vdots a_1$, i.e. $a_2 = 4a_1/3$ by (1),
- or $4a_2 - a_0 \vdots a_1$, i.e. $a_2 = a_1 + a_0/4$ by (1),
- or $3a_2 - a_0 \vdots a_1$, i.e. $a_2 = a_1 + a_0/3$ by (1).

If $a_2 = 4a_1/3$, then $d_1 = 4a_1$, $d_2 = 16a_1/3 - a_0$, $a_1 < 3a_0/2$. Then $a_4 < 240$ by Theorem QS and Lemma 2.

If $a_2 = a_1 + a_0/3$, then $a_4 = 3a_1$, $d_1 = 3a_1 + a_0$, $d_2 = 4a_1 + a_0/3$. By Theorem WF, $a_0 \vdots a_1$, i.e. $a_0 = a_1$, which violates (3).

If $a_2 = a_1 + a_0/4$, then $d_1 = 3a_1 + 3a_0/4$, $d_2 = 4a_1$. By Theorem QS (1), $i = 0$, $12a_1 \vdots a_0$, i.e. $a_0 = 4$ or $a_0 = 12$. Hence

- either $a_0 = 4$, $a_2 = a_1 + 1$, $a_3 = 2a_1$, $a_4 = 3a_1 - 1$, $d_1 = 3a_1 + 3$, $d_2 = 4a_1$,
- or $a_0 = 12$, $a_2 = a_1 + 3$, $a_3 = 2a_1$, $a_4 = 3a_1 - 3$, $d_1 = 3a_1 + 9$, $d_2 = 4a_1$.

Both these items violate conditions of Theorem WF.

4.2.3 Subcase $a_3 = a_2 - a_0$, $a_4 = a_2 + a_1 - 2a_0$

In this subcase, $d_1 = a_2 + a_1 - a_0$, $d_2 = 2a_2 + a_1 - 2a_0$. By Theorem QS (1), $i = 2$, $a_1 = 2a_0$, and hence $a_4 = a_2$, which contradicts to (3).

4.2.4 Subcase $a_3 = a_2 + a_1 - 2a_0$, $a_4 = a_2 + 2a_1 - 3a_0$

In this subcase, $d_1 = a_2 + 2a_1 - 2a_0$, $d_2 = 2a_2 + 2a_1 - 3a_0$. By Theorem QS (1), $i = 2$,

- either $2(a_1 - a_0) \vdots a_2$, i.e. $a_2 = 2(a_1 - a_0)$,
- or $2a_1 - 3a_0 \vdots a_2$, i.e. $a_2 = 2a_1 - 3a_0$.

In both cases, $a_4 < 40$ by Theorem QS (1), $i = 1$, and Lemma 2.

4.2.5 Subcase $a_3 = 2a_2 - a_1 - a_0$, $a_4 = 3a_2 - a_1 - 2a_0$

In this subcase, $d_1 = 3a_2 - a_1 - a_0$, $d_2 = 4a_2 - a_1 - 2a_0$, $a_2 < a_1 + a_0$. By Theorem QS (1), $i = 1$,

- either $3a_2 - a_0 \vdots a_1$, i.e. $a_2 = (4a_1 + a_0)/3$ by (1),
- or $4a_2 - 2a_0 \vdots a_1$, i.e. $a_2 = 5a_1/4 + a_0/2$ by (1),
- or $3a_2 - 2a_0 \vdots a_1$, i.e. $a_2 = a_1 + 2a_0/3$ by (1).

If $a_2 = (4a_1 + a_0)/3$ or $a_2 = 5a_1/4 + a_0/2$, then $a_4 < 220$ by Theorem QS and Lemma 2.

If $a_2 = a_1 + 2a_0/3$, then $a_4 = 2a_1$, $d_1 = 2a_1 + a_0$, $d_2 = 3a_1 + 2a_0/3$. By Theorem WF, $2a_0 \vdots a_1$, i.e. $a_1 = 2a_0$, $a_2 = 8a_0/3$, $a_4 = 4a_0 \leq 12$ by Lemma 2.

4.3 Case (12)

4.3.1 Subcase $a_4 = 2a_3 - a_1$

In this subcase, $d_1 = 2a_3$, $d_2 = 2a_3 + a_2 - a_1$, $a_3 < a_0 + a_1$. By Theorem QS (1), $i = 2$,

- either $2a_3 \vdots a_2$, i.e. $a_3 = 3a_2/2$ by (1),
- or $2a_3 - a_1 \vdots a_2$, i.e. $a_3 = a_2 + a_1/2$ by (1),
- or $2a_3 - a_0 \vdots a_2$, i.e. $a_3 = a_2 + a_0/2$ by (1).

In each case, $a_4 < 135$ by Theorem QS, Theorem WF and Lemma 2.

4.3.2 Subcase $a_4 = 2a_3 - a_2$

In this subcase, $d_1 = 2a_3 + a_1 - a_2$, $d_2 = 2a_3$, $a_3 < a_0 + a_2$. By Theorem QS (1), $i = 2$,

- either $2a_3 + a_1 \vdots a_2$, i.e. $a_3 = (3a_2 - a_1)/2$ or $a_3 = 2a_2 - a_1/2$ by (1),
- or $2a_3 \vdots a_2$, i.e. $a_3 = 3a_2/2$ by (1),
- or $a_3 + a_1 \vdots a_2$, i.e. $a_3 = 2a_2 - a_1$ by (1),
- or $2a_3 + a_1 - a_0 \vdots a_2$, i.e. $a_3 = (3a_2 + a_0 - a_1)/2$ by (1).

If $a_3 = 3a_2/2$ or $a_3 = 2a_2 - a_1/2$ or $a_3 = (3a_2 + a_0 - a_1)/2$, then $a_4 < 168$ by Theorem QS and Lemma 2.

If $a_3 = 2a_2 - a_1$, then $d_1 = 3a_2 - a_1$, $d_2 = 4a_2 - 2a_1$, $a_2 < a_0 + a_1$. By Theorem QS (1), $i = 1$,

- either $3a_2 \vdots a_1$, i.e. $a_2 = \lambda a_1/3$, $\lambda \in \{4, 5\}$ by (1),
- or $4a_2 \vdots a_1$, i.e. $a_2 = \lambda a_1/4$, $\lambda \in \{5, 6, 7\}$ by (1),
- or $4a_2 - a_0 \vdots a_1$, i.e. $a_2 = (\lambda a_1 + a_0)/4$, $\lambda \in \{4, 5, 6\}$ by (1).

Theorem QS and Lemma 2 give bound $a_4 < 264$ in all cases except for

$$a_2 = a_1 + a_0/4.$$

In this case, $d_1 = 2a_1 + 3a_0/4$, $d_2 = 2a_1 + a_0$. By Theorem QS (1), $i = 0$, $8a_1 \vdots a_0$. Since $\gcd(a_0/4, a_1) = 1$ by Theorem WF, $a_0 = 4$ or $a_0 = 8$. Hence

- either $a_0 = 4$, $a_2 = a_1 + 1$, $a_3 = a_1 + 2$, $a_4 = a_1 + 3$, $d_1 = 2a_1 + 3$, $d_2 = 2a_1 + 4$,
- or $a_0 = 8$, $a_2 = a_1 + 2$, $a_3 = a_1 + 4$, $a_4 = a_1 + 6$, $d_1 = 2a_1 + 6$, $d_2 = 2a_1 + 8$, where a_1 is odd.

If $a_0 = 4$, then a condition of Theorem WF is violated. Hence

$$a_0 = 8,\ a_1 = 2t + 1,\ a_2 = 2t + 3,\ a_3 = 2t + 5,\ a_4 = 2t + 7,\ d_1 = 4t + 8,\ d_2 = 4t + 10, \quad \text{where } t \text{ is even.}$$

This solution appears in Table 1 (No. 42).

If $a_3 = (3a_2 - a_1)/2$, then $d_1 = 2a_2$, $d_2 = 3a_2 - a_1$, $a_2 < a_1 + 2a_0$. By Theorem QS (1), $i = 1$,

- either $2a_2 \vdots a_1$, i.e. $a_2 = \lambda a_1/2$, $\lambda \in \{3, 4, 5\}$ by (1),
- or $3a_2 \vdots a_1$, i.e. $a_2 = \lambda a_1/3$, $4 \le \lambda \le 8$ by (1),
- or $3a_2 - a_0 \vdots a_1$, i.e. $a_2 = (\lambda a_1 + a_0)/3$, $3 \le \lambda \le 7$ by (1).

Theorem QS and Lemma 2 give bound $a_4 < 221$ in all cases except for

$$a_2 = a_1 + a_0/3.$$

In this case, $d_1 = 2a_1 + 2a_0/3$, $d_2 = 2a_1 + a_0$. By Theorem QS (1), $i = 0$, $6a_1 \vdots a_0$, i.e. $a_0 = 6$, because $6 \mid a_0$ and $\gcd(a_0/3, a_1) = 1$ by Theorem WF. Hence

$$a_0 = 6,\ a_2 = a_1 + 2,\ a_3 = a_1 + 3,\ a_4 = a_1 + 4,\ d_1 = 2a_1 + 4,\ d_2 = 2a_1 + 6,\ \text{ where } a_1 \equiv 1 \pmod 3 \text{ is odd.}$$

This solution appears in Table 1 (No. 38).

4.3.3 Subcase $a_2 = 2a_1 - a_0$, $a_4 = 2a_3 + a_0 - a_1$

In this subcase, $a_4 < a_0 + a_3$ by (1), which implies that $a_3 < a_1$. This contradicts to (3).

4.3.4 Subcase $a_3 = 2a_2 - a_1 - a_0$, $a_4 = 3a_2 - 2a_1 - a_0$

In this subcase, $a_4 < a_0 + a_3$ by (1), which implies that $a_2 < a_0 + a_1$. Hence $a_3 < a_2$, which contradicts to (3).

This finishes the proof of the Main Theorem.

Acknowledgement

We are grateful to Beijing International Center for Mathematical Research, Simons Foundation and Peking University for support, excellent working conditions and encouraging atmosphere.

Appendix

Table 1: Infinite series of well-formed quasi-smooth codimension 2 weighted complete intersection del Pezzo surfaces, which are not intersections with linear cones.

No.	$(a_0,\ a_1,\ a_2,\ a_3,\ a_4;\ d_1,\ d_2)$	I	conditions on the parameters
1	$(a_0,\ a_1,\ (\nu a_0-1)a_1,$ $\nu a_0a_1-\frac{a_0+a_1}{2},\ (\nu a_1-1)a_0;$ $\nu a_0a_1,\ 2\nu a_0a_1-a_0-a_1)$	$\frac{a_0+a_1}{2}$	$\gcd\left(\nu a_0-1,\frac{a_1-a_0}{2}\right)=1,$ $\gcd(a_0,a_1)=1,\ a_0,a_1$ odd $\nu\geq\max(1,3-a_0)$
2	$(a_0,\ a_1,\ \left(\frac{\nu a_0-1}{2}\right)a_1,$ $\left(\frac{\nu a_1-1}{2}\right)a_0,\ \left(\frac{\nu a_0+1}{2}\right)a_1-a_0;$ $\left(\frac{\nu a_0+1}{2}\right)a_1,\ (\nu a_1-1)a_0)$	$\frac{a_0+a_1}{2}$	$\gcd\left(\frac{\nu a_0-1}{2},\frac{a_1-a_0}{2}\right)=1,$ $\gcd(a_0,a_1)=1,\ \nu,a_0,a_1$ odd $\nu\geq\max(1,4-a_0)$
3	$(2b_0,\ 2b_1,\ (\nu b_0-1)b_1+b_0,$ $\nu b_0b_1,\ (\nu b_0+1)b_1-b_0;$ $(\nu b_0+1)b_1+b_0,\ 2\nu b_0b_1)$	b_0+b_1	$\gcd(\nu,b_1-b_0)=1,$ $\gcd(b_0,b_1)=1,\ \nu,b_0,b_1$ odd $\nu\geq\max(1,4-b_0)$
4	$(a_0,\ a_1,\ \nu a_0a_1+\frac{a_0-a_1}{2},$ $\nu a_0a_1,\ \nu a_0a_1+\frac{a_1-a_0}{2};$ $\nu a_0a_1+\frac{a_0+a_1}{2},\ 2\nu a_0a_1)$	$\frac{a_0+a_1}{2}$	$\gcd\left(\nu,\frac{a_1-a_0}{2}\right)=1,$ $\gcd(a_0,a_1)=1,\ a_0,a_1$ odd $\nu\geq\max(1,3-a_0)$
5	$(a_0,\ a_1,\ \left(\frac{\nu a_1+1}{2}\right)a_0-a_1,$ $\left(\frac{\nu a_0-1}{2}\right)a_1,\ \left(\frac{\nu a_1-1}{2}\right)a_0;$ $\left(\frac{\nu a_1+1}{2}\right)a_0,\ (\nu a_0-1)a_1)$	$\frac{a_0+a_1}{2}$	$\gcd\left(\frac{\nu a_0-1}{2},\frac{a_1-a_0}{2}\right)=1,$ $\gcd(a_0,a_1)=1,\ \nu,a_0,a_1$ odd $\nu\geq\max(1,6-a_0)$
6	$(a_0,\ a_1,\ 2a_1-a_0,$ $(\nu a_0-1)(2a_1-a_0),\ \nu a_0(2a_1-a_0)-a_1;$ $(\nu a_0-1)(2a_1-a_0)+a_1,\ \nu a_0(2a_1-a_0))$	a_1	$\gcd(\nu a_0-1,a_1-a_0)=1,$ $\gcd(a_0,a_1)=1,\ a_0$ odd $\nu\geq\max(1,3-a_0)$
7	$(a_0,\ a_1,\ 2a_1-a_0,\ \left(\frac{\nu a_0-1}{2}\right)(2a_1-a_0)-a_1,$ $\left(\frac{\nu a_0-1}{2}\right)(2a_1-a_0)-a_0;$ $\left(\frac{\nu a_0-1}{2}\right)(2a_1-a_0),\ \left(\frac{\nu a_0+1}{2}\right)(2a_1-a_0)-a_1)$	a_1	$\gcd\left(\frac{\nu a_0-3}{2},a_1-a_0\right)=1,$ $\gcd(a_0,a_1)=1,\ \nu,a_0$ odd $\nu\geq\max(1,6-a_0)$
8	$(a_0,\ a_1,\ 2a_1-a_0,\ \left(\frac{\nu a_0-1}{2}\right)(2a_1-a_0),$ $\left(\frac{\nu a_0-1}{2}\right)(2a_1-a_0)+a_1-a_0;$ $\left(\frac{\nu a_0-1}{2}\right)(2a_1-a_0)+a_1,\ \left(\frac{\nu a_0+1}{2}\right)(2a_1-a_0))$	a_1	$\gcd\left(\frac{\nu a_0-1}{2},a_1-a_0\right)=1,$ $\gcd(a_0,a_1)=1,\ \nu,a_0$ odd $\nu\geq\max(1,4-a_0)$
9	$(a_0,\ a_1,\ 2a_1-a_0,$ $\nu a_0(2a_1-a_0)-a_1,\ \nu a_0(2a_1-a_0)-a_0;$ $\nu a_0(2a_1-a_0),\ (\nu a_0+1)(2a_1-a_0)-a_1)$	a_1	$\gcd(\nu a_0-1,a_1-a_0)=1,$ $\gcd(a_0,a_1)=1,\ a_0$ odd $\nu\geq\max(1,3-a_0)$

10	$(2b_0,\ 2b_1,\ (2\nu b_0-1)b_1,$ $(2\nu b_1-1)b_0,\ 2(2\nu b_0 b_1-b_0-b_1);$ $2(2\nu b_0-1)b_1,\ 2(2\nu b_1-1)b_0)$	b_0+b_1	$\gcd(b_0,b_1)=1,$ b_0,b_1 odd $\nu\geq\max(1,3-b_0)$
11	$(a_0,\ a_1,\ \left(\frac{\nu a_0-1}{2}\right)a_1,$ $\left(\frac{\nu a_1-1}{2}\right)a_0,\ \nu a_0a_1-a_0-a_1;$ $(\nu a_0-1)a_1,\ (\nu a_1-1)a_0)$	$\frac{a_0+a_1}{2}$	$\gcd(a_0,a_1)=1,$ ν,a_0,a_1 odd $\nu\geq\max(1,4-a_0)$
12	$(a_0,\ a_1,\ (\nu a_0-1)a_1,$ $\nu a_0a_1-\frac{a_0+a_1}{2},\ 2\nu a_0a_1-a_0-2a_1;$ $2(\nu a_0-1)a_1,\ 2\nu a_0a_1-a_0-a_1)$	$\frac{a_0+a_1}{2}$	$\gcd(a_0,a_1)=1,$ a_0,a_1 odd $\nu\geq\max(1,3-a_0)$
13	$(a_0,\ a_1,\ (\nu a_0-1)a_1+\frac{a_1-a_0}{2},$ $(\nu a_1-1)a_0,\ 2\nu a_0a_1-2a_0-a_1;$ $2\nu a_0a_1-a_0-a_1,\ 2(\nu a_1-1)a_0)$	$\frac{a_0+a_1}{2}$	$\gcd(a_0,a_1)=1,$ a_0,a_1 odd $\nu\geq\max(1,3-a_0)$
14	$(a_0,\ a_1,\ \left(\frac{\nu a_1-1}{2}\right)a_0-a_1,$ $\left(\frac{\nu a_0-1}{2}\right)a_1-a_0,\ \nu a_0a_1-2(a_0+a_1);$ $\nu a_0a_1-a_0-2a_1,\ \nu a_0a_1-2a_0-a_1)$	$\frac{a_0+a_1}{2}$	$\gcd(a_0,a_1)=1,$ ν,a_0,a_1 odd $\nu\geq\max(1,6-a_0)$
15	$(1,\ 1,\ t,\ t,\ 2t-1;\ 2t,\ 2t)$	1	
16	$(1,\ 2,\ 2t+1,\ 2t+1,\ 4t+1;\ 4t+2,\ 4t+3)$	1	
17	$(1,\ t,\ 2t-1,\ 2t-1,\ 3t-2;\ 3t-1,\ 4t-2)$	t	
18	$(1,\ 4t-2,\ 6t-3,\ 9t-5,\ 12t-7;\ 12t-6,\ 18t-10)$	t	
19	$(1,\ 2t-1,\ 2t-1,\ 3t-2,\ 4t-3;\ 4t-2,\ 6t-4)$	t	
20	$(1,\ 6t-1,\ 8t-2,\ 12t-3,\ 18t-5;\ 18t-4,\ 24t-6)$	$2t$	
21	$(2,\ 2,\ 2t+1,\ 2t+1,\ 4t;\ 4t+2,\ 4t+2)$	2	
22	$(2,\ 3,\ 3t,\ 3t+1,\ 3t+1;\ 3t+3,\ 6t+2)$	2	
23	$(2,\ 3,\ 3t+1,\ 3t+2,\ 6t+1;\ 6t+3,\ 6t+4)$	2	
24	$(2,\ 4,\ 2t+3,\ 2t+3,\ 4t+4;\ 4t+6,\ 4t+8)$	2	
25	$(2,\ 2t+1,\ 2t+1,\ 4t+1,\ 6t+1;\ 6t+3,\ 8t+2)$	1	
26	$(3,\ t+2,\ 2t+1,\ 2t+1,\ 3t;\ 3t+3,\ 4t+2)$	$t+2$	$t\not\equiv 1\pmod 3$
27	$(3,\ 3t,\ 3t+1,\ 3t+1,\ 3t+2;\ 6t+2,\ 6t+3)$	2	
28	$(3,\ t+2,\ t+3,\ t+3,\ 2t+3;\ 2t+6,\ 3t+6)$	2	$3\nmid t$
29	$(3,\ 3t+1,\ 3t+2,\ 6t+1,\ 9t;\ 9t+3,\ 12t+2)$	2	
30	$(3,\ 2t+1,\ 2t+1,\ 3t,\ 4t-1;\ 4t+2,\ 6t)$	$t+2$	$t\not\equiv 1\pmod 3$
31	$(4,\ 6,\ 6t+5,\ 6t+7,\ 12t+8;\ 12t+12,\ 12t+14)$	4	

32	$(4,\ 6,\ 6t+3,\ 6t+5,\ 6t+5;\ 6t+9,\ 12t+10)$	4	
33	$(4,\ 2t+3,\ 2t+5,\ 4t+6,\ 6t+7;\ 6t+11,\ 8t+12)$	2	
34	$(4,\ 2t+3,\ 2t+3,\ 2t+5,\ 4t+4;\ 4t+8,\ 6t+9)$	2	
35	$(4,\ 2t+3,\ 2t+3,\ 4t+4,\ 6t+5;\ 6t+9,\ 8t+8)$	2	
36	$(4,\ 2t+3,\ 4t+6,\ 6t+7,\ 8t+8;\ 8t+12,\ 12t+14)$	2	
37	$(4,\ 4t+1,\ 4t+2,\ 4t+3,\ 4t+3;\ 8t+4,\ 8t+6)$	3	
38	$(6,\ 6t+1,\ 6t+3,\ 6t+4,\ 6t+5;\ 12t+6,\ 12t+8)$	5	
39	$(6,\ 6t+3,\ 6t+5,\ 6t+5,\ 6t+7;\ 12t+10,\ 12t+12)$	4	
40	$(6,\ 6t+3,\ 6t+5,\ 6t+5,\ 12t+4;\ 12t+10,\ 18t+9)$	4	
41	$(7,\ 4t+6,\ 6t+9,\ 9t+10,\ 12t+11;\ 12t+18,\ 18t+20)$	$t+5$	$t \not\equiv 2 \pmod 7$
42	$(8,\ 4t+5,\ 4t+7,\ 4t+9,\ 4t+11;\ 8t+16,\ 8t+18)$	6	
43	$(8,\ 4t+5,\ 4t+7,\ 4t+9,\ 8t+6;\ 8t+14,\ 12t+15)$	6	
44	$(9,\ 3t+8,\ 3t+11,\ 6t+13,\ 9t+15;\ 9t+24,\ 12t+26)$	6	
45	$(9,\ 3t+8,\ 3t+11,\ 3t+14,\ 6t+13;\ 6t+22,\ 9t+27)$	6	

In Table 1, $a_0 < a_1$, $b_0 < b_1$, ν, t denote positive integers. Note that, when $t \geq 2$, No. 19 is a special case of No. 1 (when $a_0 = 1$, $\nu = 2$), of No. 2 (when $a_0 = 1$, $\nu = 3$), of No. 11 (when $a_0 = 1$, $\nu = 3$) and of No. 12 (when $a_0 = 1$, $\nu = 2$). No. 30 is a special case of No. 2 (when $a_0 = 3$, $\nu = 1$) and of No. 11 (when $a_0 = 3$, $\nu = 1$). No. 17 is a special case of No. 6 (when $a_0 = 1$, $\nu = 2$) and of No. 8 (when $a_0 = 1$, $\nu = 3$). No. 26 is a special case of No. 8 (when $a_0 = 3$, $\nu = 1$).

Table 2 below lists well-formed quasi-smooth codimension 2 weighted complete intersection del Pezzo surfaces, which are not intersections with linear cones, with $a_4 \leq 500$, $d_2 \leq 1000$, which do not appear in Table 1. It was obtained by running a computer program, whose code is available from the author upon request. According to the Main Theorem, Table 1 and Table 2 together list all well-formed quasi-smooth codimension 2 weighted complete intersection del Pezzo surfaces, which are not intersections with linear cones.

Table 2: Sporadic well-formed quasi-smooth codimension 2 weighted complete intersection del Pezzo surfaces, which are not intersections with linear cones.

$(a_0,\ a_1,\ a_2,\ a_3,\ a_4;\ d_1,\ d_2)$	$(a_0,\ a_1,\ a_2,\ a_3,\ a_4;\ d_1,\ d_2)$	$(a_0,\ a_1,\ a_2,\ a_3,\ a_4;\ d_1,\ d_2)$
(13, 22, 55, 76, 97; 110, 152)	(9, 19, 24, 31, 53; 62, 72)	(1, 7, 11, 17, 27; 28, 34)
(11, 27, 36, 62, 97; 108, 124)	(14, 19, 25, 32, 45; 64, 70)	(5, 7, 10, 14, 23; 28, 30)
(13, 18, 45, 61, 77; 90, 122)	(10, 17, 25, 34, 43; 60, 68)	(2, 7, 10, 13, 18; 20, 28)
(11, 29, 39, 49, 67; 78, 116)	(11, 17, 24, 31, 37; 48, 68)	(6, 7, 9, 11, 14; 18, 28)

(11, 29, 38, 48, 85; 96, 114)	(11, 14, 21, 33, 52; 63, 66)	(5, 8, 9, 12, 19; 24, 27)
(13, 23, 35, 57, 79; 92, 114)	(13, 14, 23, 33, 43; 56, 66)	(2, 7, 8, 13, 19; 21, 26)
(13, 23, 34, 56, 89; 102, 112)	(13, 14, 23, 32, 33; 46, 65)	(1, 5, 9, 13, 17; 18, 26)
(13, 23, 35, 47, 57; 70, 104)	(11, 15, 20, 32, 49; 60, 64)	(5, 6, 9, 13, 13; 18, 26)
(11, 25, 34, 43, 57; 68, 100)	(11, 17, 24, 31, 38; 55, 62)	(6, 8, 9, 11, 13; 22, 24)
(11, 29, 39, 49, 59; 88, 98)	(10, 13, 25, 31, 37; 50, 62)	(1, 5, 8, 12, 19; 20, 24)
(13, 20, 31, 49, 67; 80, 98)	(11, 17, 20, 27, 43; 54, 60)	(5, 7, 8, 11, 14; 21, 22)
(11, 25, 32, 41, 71; 82, 96)	(9, 15, 23, 23, 37; 46, 60)	(2, 5, 8, 11, 14; 16, 22)
(13, 20, 29, 47, 74; 87, 94)	(13, 14, 19, 29, 44; 57, 58)	(3, 7, 8, 9, 13; 16, 21)
(11, 21, 28, 47, 73; 84, 94)	(14, 15, 19, 26, 37; 52, 56)	(4, 5, 7, 10, 13; 18, 20)
(13, 14, 35, 46, 57; 70, 92)	(9, 15, 23, 23, 31; 46, 54)	(1, 4, 7, 10, 13; 14, 20)
(13, 20, 31, 42, 49; 62, 91)	(13, 14, 19, 23, 29; 42, 52)	(3, 5, 6, 8, 13; 16, 18)
(9, 23, 30, 38, 67; 76, 90)	(11, 13, 19, 25, 27; 38, 52)	(3, 5, 7, 9, 11; 16, 18)
(11, 18, 27, 44, 61; 72, 88)	(11, 17, 20, 24, 27; 44, 51)	(2, 5, 6, 9, 13; 15, 18)
(11, 25, 34, 43, 52; 77, 86)	(5, 14, 17, 21, 37; 42, 51)	(3, 5, 7, 9, 11; 14, 18)
(10, 19, 35, 43, 51; 70, 86)	(11, 13, 19, 25, 31; 44, 50)	(1, 4, 5, 7, 11; 12, 15)
(14, 17, 29, 41, 44; 58, 85)	(9, 12, 19, 19, 29; 38, 48)	(4, 5, 6, 7, 8; 12, 14)
(11, 21, 29, 37, 47; 58, 84)	(1, 9, 15, 23, 23; 24, 46)	(3, 4, 6, 7, 8; 12, 14)
(14, 17, 29, 41, 53; 70, 82)	(9, 12, 19, 19, 26; 38, 45)	(3, 4, 5, 6, 7; 11, 12)
(13, 17, 27, 41, 55; 68, 82)	(9, 10, 15, 22, 23; 32, 45)	(3, 4, 5, 6, 7; 10, 12)
(14, 17, 27, 39, 64; 78, 81)	(10, 11, 15, 22, 29; 40, 44)	(2, 3, 5, 6, 7; 10, 12)
(11, 21, 26, 34, 57; 68, 78)	(11, 13, 14, 20, 29; 40, 42)	(3, 3, 5, 5, 7; 10, 12)
(13, 20, 29, 31, 47; 60, 78)	(5, 9, 12, 20, 31; 36, 40)	(2, 3, 4, 5, 5; 8, 10)
(13, 17, 27, 37, 41; 54, 78)	(1, 8, 13, 19, 31; 32, 39)	(1, 3, 3, 5, 5; 6, 10)
(13, 17, 24, 38, 59; 72, 76)	(2, 9, 12, 19, 19; 21, 38)	(2, 2, 3, 3, 3; 6, 6)
(11, 25, 32, 34, 41; 66, 75)	(9, 11, 12, 17, 25; 34, 36)	(1, 2, 2, 3, 3; 4, 6)
(11, 21, 29, 37, 45; 66, 74)	(1, 7, 12, 17, 23; 24, 35)	

References

[1] C. Boyer, K. Galicki, and M. Nakamaye, *On the geometry of Sasakian-Einstein* 5*-manifolds*, Mathematische Annalen **325** (2003), no. 3, 485–524.

[2] I. Cheltsov and C. Shramov, *Del Pezzo zoo*, Experimental Mathematics **22** (2013), no. 3, 313–326.

[3] J.-J. Chen, J. A. Chen, and M. Chen, *On quasismooth weighted complete intersections*, Journal of Algebraic Geometry **20** (2011), no. 2, 239–262.

[4] I. Dolgachev, *Weighted projective varieties*, Group actions and vector fields (Vancouver, B.C., 1981), Lecture Notes in Mathematics, vol. 956, Springer, Berlin, 1982, pp. 34–71.

[5] A. R. Iano-Fletcher, *Working with weighted complete intersections*, Explicit birational geometry of 3-folds, London Mathematical Society Lecture Note Series, vol. 281, Cambridge University Press, Cambridge, 2000, pp. 101–173.

[6] J. M. Johnson and J. Kollár, *Kähler-Einstein metrics on log del Pezzo surfaces in weighted projective* 3*-spaces*, Université de Grenoble. Annales de l'Institut Fourier **51** (2001), no. 1, 69–79.

[7] I.-K. Kim and J. Park, *Log canonical thresholds of complete intersection log del Pezzo surfaces*, Proceedings of the Edinburgh Mathematical Society. Series II **58** (2015), no. 2, 445–483.